

Annual Report

2017

Volume 12

Kagawa School of
Pharmaceutical
Sciences

Tokushima Bunri University

Kagawa School of Pharmaceutical Sciences
Tokushima Bunri University

Shido 1314-1, Sanuki
Kagawa 769-2193, JAPAN

Telephone: +81-87-899-7100

Facsimile: +81-87-894-0181

URL: <http://kp.bunri-u.ac.jp/index.html>

Preface to the tenth issue of *The Annual Report*

The Faculty of Pharmaceutical Sciences at the Kagawa Campus was founded in April, 2004. Our purpose is to educate students to become professional pharmacists and basic scientists with sufficient knowledge of chemistry, biology, and pharmaceutical thought to practice with humanity and a high sense of ethics and social responsibility. In April, 2005, the Kagawa School of Pharmaceutical Sciences was established with masters and doctoral programs. The undergraduate course of the Kagawa School of Pharmaceutical Sciences was reorganized in April, 2006, into two departments, the Department of Pharmacy (with a six-year training program) and Pharmaceutical Sciences (with a four-year training program). The mission of the Department of Pharmacy (the six-year program) is to train highly qualified pharmacists who are knowledgeable about advanced medical treatment, and who also can contribute to the prevention and treatment of diseases. Society's view of pharmacy education has changed, not only in medical fields but also in other fields, to the fostering of a healthy and safe society. In order to fulfill this mandate, the Department of Pharmaceutical Sciences (the four-year program) educates students to become key members of society who can play an active part in all social fields with their broad background in pharmacy, and who can become outstanding life scientists by making signal contributions to basic science. But Kagawa School of Pharmaceutical Sciences has turned into only the program of the Department of Pharmacy in April, 2012, in order to match the needs of the region. Our faculty, which has cutting-edge equipment, succeeds the principles at the foundation and has trained professionals with a research mind who can respond the social demands, and a variety of human resources who can contribute to the promotion of health of the people.

Guided by these principles, we laid the foundation for the Kagawa School of Pharmaceutical Sciences and dedicated ourselves to this effort. We established Institute of Neuroscience in the Kagawa School of Pharmaceutical Sciences, and the Institute of Neuroscience was reorganized and has now been expanded into four divisions in April, 2007. We published the first issue of *The Annual Report of the Kagawa School of Pharmaceutical Sciences*, highlighting our educational, research, management, and philanthropic achievements, by looking back upon the past three years of our activities, in 2006. We are now publishing the eleventh issue of the *Annual Report* by looking back upon the past five years of our activities. I would like all of the staff to utilize these issues to perform a self-assessment and to look forward to the future development of their activities. In addition, I expect that this issue will assist our staff in expanding both intramural and extramural interdisciplinary exchanges.

The Kagawa School of Pharmaceutical Sciences is always making progress on the road toward realizing our goals. I would be very grateful if all the people who read this annual report would kindly give me their opinions and/or comments with a view toward continuing to improve the Kagawa School of Pharmaceutical Sciences.

Hiroshi Miyazawa, Ph.D., Dean
Kagawa School of Pharmaceutical Sciences
Tokushima Bunri University

Contents

Preface to the eleventh issue of *The Annual Report*

Short History of <i>The Kagawa School of Pharmaceutical Sciences</i>	• • • • • 135
---	---------------

Reports

Laboratory for Analytical Chemistry	• • • • • 137
Laboratory of Biophysics	• • • • • 143
Laboratory of Pharmacognosy and Natural Products Chemistry	• • • • • 147
Laboratory of Medicinal Chemistry	• • • • • 151
Laboratory for Molecular Biology	• • • • • 153
Laboratory of Immunology	• • • • • 159
Laboratory of Pharmaceutical Health Sciences	• • • • • 161
Laboratory of Neuropharmacology	• • • • • 163
Laboratory of Pathological Physiology	• • • • • 165
Laboratory for Pharmacotherapy and Experimental Neurology	• • • • • 167
Laboratory of Pharmacokinetics and Pharmacodynamics	• • • • • 171
Laboratory of Pharmaceutics	• • • • • 175
Laboratory of Pharmaceutical Health Care and Sciences	• • • • • 177
Laboratory of Pharmaceutical Education	• • • • • 179
Center for Instrumental Analysis	• • • • • 183
Laboratory for Neural Circuit Systems , Institute of Neuroscience	• • • • • 185

A Short History of The Kagawa School of Pharmaceutical Sciences

- 1895 Sai Murasaki founds a private vocational school in the city of Tokushima.
- 1966 Tokushima Women's University is founded.
Faculty of Home Economics admits its first students.
- 1968 Faculty of Music admits its first students.
- 1972 Tokushima Women's University is renamed Tokushima Bunri University.
Faculty of Pharmaceutical Sciences admits its first students.
- 1983 The Kagawa Campus of Tokushima Bunri University opens.
Faculty of Literature admits its first students on the Kagawa campus.
- 1989 Faculty of Engineering admits its first students on the Kagawa campus.
- 1995 The 100th anniversary of the founding of the Murasaki Gakuen.
- 2000 Faculty of Policy Studies admits its first students.
- 2004 Faculty of Pharmaceutical Sciences at Kagawa Campus admits its first students on the Kagawa campus; the Department of Pharmaceutical Technology opens 10 laboratories.
The rooms for teaching staff open on the fourth floor of the Research and Media Library.
The office of the Faculty of Pharmaceutical Sciences at Kagawa Campus opens on the first floor of the Lecture Building.
- 2005 The Kagawa School of Pharmaceutical Sciences is established on the Kagawa campus.
Masters and doctoral programs begin at the Kagawa School of Pharmaceutical Sciences.
Pharmaceutical Sciences Research and Laboratory Buildings are completed.
The Center for Instrumental Analysis opens in the Pharmaceutical Sciences Research Building.
The Center for Radioisotope and Laboratory Animals opens in the Pharmaceutical Sciences Laboratory Building.
The Medicinal Plant Garden is established.
The 110th anniversary of the founding of the Murasaki Gakuen.
- 2006 The six-year undergraduate program in the Department of Pharmacy, Kagawa School of Pharmaceutical Sciences accepts its first students.
The four-year undergraduate program in the Department of Pharmaceutical Sciences, Kagawa School of Pharmaceutical Sciences accepts its first students.
A mock pharmacy opens in the Lecture Building for student training.
The construction of eighteen laboratories and one research institute for the Kagawa School of Pharmaceutical Sciences is completed.
SENKA Endowed Chair for "Practical Drug Discovery and Development" was established.
- 2007 The research institute (Institute of Neuroscience) was reorganized and expanded into four divisions.
- 2012 The four-year Graduate School of Pharmaceutical Sciences (Department of Pharmaceutical Sciences Doctoral program) was installed.

- 2014 The 10th anniversary of the foundation of Kagawa School of Pharmaceutical Sciences
- 2015 The 120th anniversary of the founding of the Murasaki Gakuen.



Laboratory for Analytical Chemistry

Staff

Professor: Kentaro Yamaguchi, Ph. D (Apr. 2004)

Educational History:

Graduated from University of

Electrocommunications in Mar. 1975

Ph. D (Tokyo University)

Latest Work Record:

Associate Professor in Chiba University

Associate Professor: Masahide Tominaga, Ph.D.

Educational History:

D. Eng. University of Tokyo, 2000

Lecturer: Masatoshi Kawahata, Ph. D

Educational History:

Graduated from Graduate School of Tokushima

Bunri University in Mar. 2006

Research

Laboratory of Analytical chemistry has designated that it contributes to pharmaceutical technology with analyzing molecular structures, physical properties, chemical reactivity, and functions. The connection between dynamic behavior of biomolecules and its reactivity-function relationship as well as the detection of unstable reaction intermediate in solution are our major subjects. We have been engaged in the study to produce new analysis method, which reveals the generating mechanism of the outstanding functionalities and the physical properties of the materials in solution based on microanalysis in the atomic resolution. We also develop the analytical and chemical technique in solution that is based on quantum mechanics and statistical mechanics. These results might be applied to pharmaceutical organic synthesis.

Development in the field of mass spectrometry:

In the laboratory, we have succeeded in the development of the 'Cold-Spray Ionization (CSI)' method. CSI is designed for mass spectrometric detection of labile organic species. It may be an appropriate method to analyze in solution the structures of biomolecular complexes, labile organic species including Grignard reagents, asymmetric catalysis, and supermolecules. The method, a variant of Electrospray ionization (ESI)-MS, operates at low temperature, allow simple and precise characterization of labile non-covalent complexes that are difficult or impossible to observe by conventional MS techniques, including fast atom bombardment (FAB), and matrix assisted laser desorption ionization (MALDI), as well as ESI.

Biopolymer analysis:

The behavior of important in vivo molecule such as protein, nucleic acid, and lipid, will be made clear by using our newly developed method CSI mass spectrometry. The structure of multistranded DNA of duplex, triplex and quadruplex DNA have been examined by electrospray ionization (ESI) MS. However, non-covalent complexes of multiply stranded DNA are difficult to observe by conventional methods, because of their low melting temperature (T_m). The characterization of triple- and

quadruple-stranded oligodeoxynucleotides was carried out by means of CSI-MS. In consequence, it is proved that DNA has been made clear to take hyper-stranded structures combining various multiply stranded helical components.

Analysis of organic reaction mechanism:

A new CSI mass spectrometric procedure RTS (reaction tracking system), which can trace the time-dependent simultaneous change of the raw materials, products and inter mediates in an organic reaction under CSI condition has been developed. It would be expected that behavior of each component becomes clear and that the appropriate design become possible to make the reaction more efficiently.

Observation of the giant molecules by means of mass spectrometry:

Mass Spectrometry (MS) has been developed and adopted to wide variety of analytical chemistry in recent years.

Although MS was basically developed for high molecular weight substances in the field of biochemistry, the measurement of huge molecules over 10k Da is still very difficult. This is caused by the ionizing problems, stability of the compounds and the existence of various impurities.

We develop some new techniques to overcome these problems by using newly equipped FT-ICR mass spectrometer.

Construction of Adamantane-Based Macrocycles and Cages:

Synthetic macrocyclic compounds are considerably important receptor molecules, and they provide an opportunity to interact with various guest molecules by the binding sites of multiple functional groups and well-defined cavities within their frameworks. Adamantane derivatives are unique and specific compounds, because they are mechanically rigid and conformationally well-defined. However, the design and synthesis of adamantane-based host molecules and their applications in host-guest systems remain largely unexplored. We reported the construction and structural analysis of various types of nano-sized adamantane-based macrocycles and cages bearing rectangle, square, and spherical shapes. Their macrocycles encapsulate electron-poor guest molecule, 1,3,5-trinitrobenzene via charge-transfer interactions. Further, adamantane-based macrocycles showed the generation of unique molecular networks in the solid state. The adamantane-based macrocycles formed tubular structures with channels, which were assembled into three-dimensional networks through multiple intermolecular interactions.

Self-assembled Nano-scale Structures Using Macrocyclic Framework as Molecular Block:

Recently, several functionalized macrocycles were utilized as building blocks for self-assembled nanostructures including vesicles, tubes, fibers, and others. We synthesized a series of oxacyclophanes, azacyclophanes, and salen-based macrocycles bearing adamantane parts. By utilizing the specific properties of spherical shape, bulky skeleton, and lipophilicity of the adamantane derivatives, these adamantane-derived macrocycles were induced into hollow spherical aggregates with a multilayer membrane in organic and aqueous solutions. The hollow spheres were induced into fibrous and/or network aggregates, which were eventually transformed into single crystals. These results afforded the evidences for a morphological change and phase transition occurring from the solution into a solid.

Publications (2013~2018)

[Original papers]

2018

1. Hayashi, Y.; Ohara, K.; Taki, R.; Saeki, T.; *Yamaguchi, K. (2018). Combined analysis of 1,3-benzodioxoles by crystalline sponge X-ray crystallography and laser desorption ionization mass spectrometry. *Analyst*, 2018, 143, 1475-1481.
2. Yamada, M.; Matsumura, M.; Takino, F.; Murata, Y., Kurata, Y.; Kawahata, M.; Yamaguchi, K.; Kakusawa, N.; *Yasuike, S. (2018). Synthesis of Fully Functionalized 5-Selanyl-1,2,3-triazoles: Copper-Catalysed Three-Component Reaction of Ethynylstibanes, Organic Azides, and Diaryl Diselenides. *Eur. J. Org. Chem.*, 2018, 2, 170-177.
3. Hirao, T.; Kim, D-S.; Chi, X.; Lynch, V-M., Ohara, K.; *Park, J-S.; *Yamaguchi, K.; *Sessler, J-L. (2018). Control over multiple molecular states with directional changes driven by molecular recognition. *Nat. Commun.*, 2018, 9, 823.
4. Ning, Y.; Kawahata, M.; Yamaguchi, K.; Otani, Y.; *Ohwada, T. (2018). Synthesis, Structure and N-N Bonding Character of 1,1-Disubstituted Indazolium Hexafluorophosphate. *Chem. Commun.*, 2018, 54, 1881-1884.
5. *Tominaga, M.; Ando, H.; Ohara, K.; Itho, T.; Yamaguchi, K. (2018). Crystal Formation of a Coordination Cage through Spherical Particles Derived from a Tripodal Ligand Containing Adamantane Moiety and Silver(I) Salt. *Chem. Lett.*, 2018, 47, 315-317.

2017

1. Suzuki, N.; Kawahata, M.; Yamaguchi, K.; Suzuki, T.; Tomono, K.; Fukami, T. (2017). Comparison of the relative stability of pharmaceutical cocrystals consisting of paracetamol and dicarboxylic acids. *Drug Dev. Ind. Pharm.*, 2017, 44, 582-589.

2. *Kawahata, M.; Tominaga, M.; Maekawa, Y.; *Yamaguchi, K. (2017). Preparation and crystal structures of charge-transfer complexes of acyclic host molecules bearing pyrogallol derivatives with paraquat. *CrystEngComm*, 2017, 19, 7229-7235.
3. *Katagiri, K.; Matsuo, N.; Kawahata, M.; Masu, H.; Yamaguchi, K. (2017). Aqua coordination to attenuate the luminescence properties of europium(III)-phosphine oxide porous coordination polymers. *New J. Chem.*, 2017, 41, 8305-8311.
4. *Danjo, H.; Hamaguchi, M.; Asai, K.; Nakatani, M.; Kawanishi, H.; Kawahata, M.; Yamaguchi, K. (2017). Proton-Induced Assembly-Disassembly Modulation of Spiroborate Twin-Bowl Polymers Bearing Pyridyl Groups. *Macromolecules*, 2017, 50, 8028-8032.
5. Terazawa, S.; Uemura, Y.; Koyama, Y.; Kawakami, S.; Sugimoto, S.; Matsunami, K.; *Otsuka, H.; Shinzato, T.; Kawahata, M.; Yamaguchi, K. (2017). Microtropins Q-W, *ent*-Labdane Glucosides: Microtropiosides G-I, Ursane-Type Triterpene Diglucoside and Flavonol Glycoside from the Leaves of *Microtropis japonica*. *Chem. Pharm. Bull.*, 2017, 65, 930-939.
6. *Tominaga, M.; Takahashi, E.; Ukai, H.; Ohara, K.; Itho, T.; Yamaguchi, K. (2017). Solvent-Dependent Self-Assembly and Crystal Structures of a Salen-Based Macrocycle. *Org. Lett.*, 2017, 19(7), 1508-1511.
7. Ning, Y.; Fukuda, T.; Ikeda, H.; Otani, Y.; Kawahata, M.; Yamaguchi, K.; *Ohwada, T. (2017). Revisiting secondary interactions in neighboring group participation, exemplified by reactivity changes of iminylium intermediates. *Org. Biomol. Chem.*, 2017, 15, 1381-1392.

2016

1. *Tominaga, M.; Kawaguchi, T.; Ohara, K.; Yamaguchi, K.; Katagiri, K.; Itho, T.; *Azumaya, I. (2016). Vesicle Formation of Three-dimensional Trinuclear Silver(I) Complexes Built from Tris-NHC Ligands Bearing Long Alkyl Chains. *Chem. Lett.*, 2016, 45, 1201-1203.
2. *Tominaga, M.; Noda, A.; Ohara, K.; Yamaguchi, K.; Itho, T. (2016). Synthesis, Hollow Spherical Aggregation, and Crystallization of an Adamantane-derived Azacyclophane Containing Triazine Rings. *Chem. Lett.*, 2016, 45, 733-775.
3. *Sasaki, M.; Ando, M.; Kawahata, M.; Yamaguchi, K.; Takeda, K. (2016). Spontaneous Oxygenation of Siloxy-*N*-silylketenimines to α -Ketoamides. *Org. Lett.*, 2016, 18, 1598-1601.



4. *Ohara, K.; Tominaga, M.; Masu, H.; Azumaya, I.; *Yamaguchi, K. (2016). Adamantane-based Bidendate Metal Complexes in Crystalline and Solution State. *Anal. Sci.*, 2016, 32(12), 1347-1352.
5. Matsumura, M.; Sakata, Y.; Iwase, A.; Kawahata, M.; Kitamura, Y.; Murata, Y.; Kakusawa, N.; Yamaguchi, K.; Yasuike, S. (2016). Copper-catalyzed tandem cyclization of 2-(2-iodophenyl)imidazo[1,2-*a*]pyridine derivatives with selenium: Synthesis of benzo[*b*]selenophenefused imidazo[1,2-*a*]pyridines. *Tetrahedron Lett.*, 2016, 57, 5484-5488.
6. Kawakami, S.; Inagaki, M.; Matsunami, K.; *Otsuka, H.; Kawahata, M.; Yamaguchi, K. (2016). Crotofolane-Type Diterpenoids, Crotoascarins L-Q, and a Rearranged Crotofolane-Type Diterpenoid, Neocrotoascarin, from the Stems of *Croton cascarilloides*. *Chem. Pharm. Bull.*, 2016, 64, 1492-1498.
7. Wang, D-Y.; Kawahata, M.; Yang, Z-K.; Miyamoto, K.; Komagawa, S.; Yamaguchi, K.; Wang, C.; Uchiyama, M. (2016). Stille coupling via C-N bond cleavage. *Nat. Commun.*, 2016, 7, 12937.
8. *Kawahata, M.; Komagawa, S.; Ohara, K.; Fujita, M.; *Yamaguchi, K. (2016). High-resolution X-ray structure of methyl salicylate, a time-honored oily medicinal drug, solved by crystalline sponge method. *Tetrahedron Lett.*, 2016, 57, 4633-4636.
9. *Iwatsuki, S.; Ichiyama, A.; Tanooka, S.; Toyama, M.; Katagiri, K.; Kawahata, M.; Yamaguchi, K.; Danjo, H.; *Chayama, K. (2016). Coordination structure and extraction behavior of a silver ion with *N*-substituted-9-aza-3,6,12,15-tetrahydroheptadecanes: significant effect of Ph-C-N framework on the extractability. *Dalton Trans.*, 2016, 45, 12548-12558.
10. Masuda, T.; Arase, J.; Inagaki, Y.; Kawahata, M.; Yamaguchi, K.; Ohhara, T.; Nakao, A.; Momma, H.; Kwon, E.; *Setaka, W. (2016). Molecular Gyrotops with a Five-Membered Heteroaromatic Ring: Synthesis, Temperature-Dependent Orientation of Dipolar Rotors inside the Crystal, and its Birefringence Change. *Cryst. Growth Des.*, 2016, 16, 4392-4401.
11. Sawada, T.; Yamagami, M.; Ohara, K.; Yamaguchi, K.; *Fujita, M. (2016). Peptide [4]Catenane by Folding and Assembly. *Angew. Chem. Int. Ed.*, 2016, 55, 4519-4522.
12. Wang, S.; Taniguchi, T.; Monde, K.; Kawahata, M.; Yamaguchi, K.; *Otani, Y.; *Ohwada, T. (2016). Hydrogen bonding to carbonyl oxygen of nitrogen-pyramidalized amide - detection of pyramidalization direction preference by vibrational circular dichroism spectroscopy. *Chem. Commun.*, 52, 4018-4021.
13. Taro, A.; Niki, A.; Komagawa, S.; Arimitsu, K.; Uchimoto, H.; *Kawasaki, I.; Yamaguchi, K.; *Nishide, K. (2016). A highly regio- and stereoselective selenoxide elimination of ,2-bis[4-(trimethylsilyl)phenylseleno]alkanes to give (*E*)-alkenyl selenoxides and its mechanistic study. *ChemistrySelect*, 2, 189-194.
14. Ishizuka, T.; Watanabe, A.; Kotani, H.; Hong, D.; Satonaka, K.; Wada, T.; Shiota, Y.; Yoshizawa, K.; Ohara, K.; Yamaguchi, K.; Kato, S.; *Fukuzumi, S.; *Kojima, T. (2016). Homogeneous Photocatalytic Water Oxidation with a Dinuclear Co^{III}-Pyridylmethylamine Complex. *Inorg. Chem.*, 55, 1154-1164.
15. Wang, S.; Sawada, T.; Ohara, K.; Yamaguchi, K.; *Fujita, M. (2016). Capsule-Capsule Conversion by Guest Encapsulation. *Angew. Chem. Int. Ed.*, 55, 2063-2066.
16. *Tominaga, M.; Kawaguchi, T.; Ohara, K.; Yamaguchi, K.; Masu, H.; *Azumaya, I. (2016). Synthesis and crystal structures of twisted three-dimensional assemblies of adamantane-bridged tris-NHC ligands and Ag^I. *CrystEngComm*, 18, 266-273.
- ### 2015
1. Kawakami, S.; Matsunami, K.; *Otsuka, H.; Inagaki, M.; Takeda, Y.; Kawahata, M.; Yamaguchi, K. (2015). Crotoascarins I-K: Crotofolane-Type Diterpenoids, Crotoascarin γ , Isocrotofolane Glucoside and Phenolic Glycoside from the Leaves of *Croton cascarilloides*. *Chem. Pharm. Bull.*, 63, 1047-1054.
2. Kotomori, Y.; *Sasaki, M.; Kawahata, M.; Yamaguchi, K.; Takeda, K. (2015). Stereochemical Course of Deprotonation-Acylation of *N*-Boc- and *N*-Carbamoyl-2-cyano-6-methylpiperidines. *J. Org. Chem.*, 80(21), 11013-11020.
3. Nishiyama, Y.; Inagaki, Y.; Yamaguchi, K.; *Setaka, W. (2015). 1,4-Naphthalenediyl-Bridged Molecular Gyrotops: Rotation of the Rotor and Fluorescence in Solution. *J. Org. Chem.*, 80, 9959-9966.
4. Shionari, H.; Inagaki, Y.; Yamaguchi, K.; *Setaka, W. (2015). A pyrene-bridged macrocage showing no excimer fluorescence. *Org. Biomol. Chem.*, 13, 10511-10516.
5. *Ikeda, A.; Iwata, N.; Hino, S.; Mae, T.; Tsuchiya, Y.; Sugikawa, K.; Hirao, T.; Haino, T.; Ohara, K.; Yamaguchi, K. (2015). Liposome collapse resulting from an allosteric interaction between 2,6-dimethyl- β -cyclodextrins and lipids. *RSC Adv.*, 5, 77746-77754.
6. Uemura, Y.; Iwami, M.; Kawakami, S.; Sugimoto, S.; Matsunami, K.; *Otsuka, H.; Shinzato, T.; Kawahata, M.; Yamaguchi, K. (2015). Megastigmane Glucosides and

- Megastigmanes from the Leaves of *Meliosma lepidota* ssp. *squamulata*. Chem. Pharm. Bull., 63, 608-616.
7. *Katagiri, K.; Komagawa, S.; Uchiyama, M.; Yamaguchi, K.; *Azumaya, I. (2015). Control of a Chiral Property of a Calix[3]aramide: The Racemization Suppressed by Intramolecular Cyclic Hydrogen Bonds and DMSO-H₂O System-Induced Spontaneous Resolution. Org. Lett., 17, 3650-3653.
8. Tarao, A.; Tabuchi, Y.; Sugimoto, E.; Ikeda, M.; Uchimoto, H.; Arimitsu, K.; Kimura, H.; Kawasaki, I.; Kawahata, M.; Yamaguchi, K.; *Nishide, K. (2015). A mechanistic investigation of anti-elimination in (Z)-1,2-bis(arylseleno)-1-alkenes and their sulfur analogs. Org. Biomol. Chem., 13, 5964-5971.
9. *Danjo, H.; Kidena, Y.; Kawahata, M.; Sato, H.; Katagiri, K.; Miyazawa, T.; Yamaguchi, K. (2015). Multilayered Inclusion Nanocycles of Anionic Spiroborates. Org. Lett., 17, 2466-2469.
10. *Danjo, H.; Hashimoto, Y.; Kidena, Y.; Nogamine, A.; Katagiri, K.; Kawahata, M.; Miyazawa, T.; Yamaguchi, K. (2015). Nestable Tetrakis(spiroborate) Nanocycles. Org. Lett., 17, 2154-2157.
11. Ohara, K.; Nakai, A.; Yamaguchi, K. (2015). Laser desorption ionization of stilbenes in crystalline sponge. Eur. J. Mass Spectrom., 21, 413-421.
12. *Sasaki, M.; Kondo, Y.; Moto-ishi, T.; Kawahata, M.; Yamaguchi, K.; Takeda, K. (2015). Enantioselective Synthesis of Allenylenol Silyl Ethers via Chiral Lithium Amide Mediated Reduction of Ynenoyl Silanes and Their Diels-Alder Reactions. Org. Lett., 17, 1280-1283.
13. Danjo, H.; *Nakagawa, T.; Katagiri, K.; Kawahata, M.; Yoshigai, S.; Miyazawa, T.; Yamaguchi, K. (2015). Formation of Lanthanide(III)-Containing Metallosupramolecular Arrays Induced by Tris(spiroborate) Twin Bowl. Cryst. Growth. Des., 15, 384-389.
14. Katagiri, K.; Tohaya, T.; Shirai, R.; Kato, T.; Masu, H.; Tominaga, M.; Azumaya, I. (2015). Folded-to-unfolded structural switching of a macrocyclic aromatic hexamide based on conformation changes in the amide groups induced by *N*-alkylation and dealkylation reactions. J. Mol. Struct. 1082, 23-28.
15. Tominaga, M., Kunitomi, M., Katagiri, K., Itoh, T. (2015). Adamantane-Based Oxacyclophanes Containing Pyrazines: Synthesis, Crystal Structure, and Self-Assembly Behavior. Org. Lett., 17, 786-789.
- Meerwein-Ponndorf-Verley-Type Reduction of α -Silylimines by a Chiral Lithium Amide. J. Org. Chem., 79(8), 3601-3609.
2. *Tominaga, M.; Yoneta, T.; Ohara, K.; Yamaguchi, K.; Itoh, T.; Minamoto, C.; *Azumaya, I. (2014). Self-Assembly of a Tetrapodal Adamantane with Carbazole Branches into Hollow Spherical Aggregates in Organic Media. Org. Lett., 16, 4622-4625.
3. *Tominaga, M.; Ukai, H.; Katagiri, K.; Ohara, K.; Yamaguchi, K.; *Azumaya, I. (2014). Tubular Structures Bearing Channels in Organic Crystals Composed of Adamantane-Based Macrocycles. Tetrahedron, 70, 2576-2581.
4. *Tominaga, M.; Iekushi, A.; Katagiri, K.; Ohara, K.; Yamaguchi, K.; *Azumaya, I. (2014). Channel-Dependent Conformations of Single-Strand Polymers in Organic Networks Composed of Tetrapodal Adamantanes with *N*-heterocyclic Moieties. Tetrahedron Lett., 55, 5789-5792.
5. Maeno, Y.; *Fukami, T.; Kawahata, M.; Yamaguchi, K.; Tagami, T.; Ozeki, T.; Suzuki, T.; Tomono, K. (2014). Novel Pharmaceutical Cocrystal Consisting of Paracetamol and Trimethylglycine, a new Promising Cocrystal Former. Int. J. Pharm., 473, 179-186.
6. Mu, Y-J.; Yu, L-N.; *Jiang, X-F.; *Yu, S-Y.; Yamaguchi, K. (2014). Self-Assembly of an Organo-Palladium Molecular Basket that Encapsulates Cobaltcarborane Anion in Water. Inorg. Chem. Commun., 44, 119-123.
7. *Setaka, W.; Inoue, K.; Higa, S.; Yoshigai, S.; Kono, H.; Yamaguchi, K. (2014). Synthesis of Crystalline Molecular Gyrotops and Phenylene Rotation inside the Cage. J. Org. Chem., 79, 8288-8295.
8. *Tominaga, M.; Ohara, K.; Yamaguchi, K.; *Azumaya, I. (2014). Hollow Sphere Formation from a Three-Dimensional Structure Composed of an Adamantane-Based Cage. J. Org. Chem., 79, 6738-6742.
9. Wang, S.; *Otani, Y.; Liu, X.; Kawahata, M.; Yamaguchi, K.; *Ohwada, T. (2014). Robust trans-Amide Helical Structure of Oligomers of Bicyclic Mimics of β -Proline: Impact of Positional Switching of Bridgehead Substituent on Amide *cis-trans* Equilibrium. J. Org. Chem., 79, 5287-5300.
10. Inagaki, Y.; Yamaguchi, K.; *Setaka, W. (2014). A Crystalline Molecular Gyrotop with Germanium Junctions Between a Phenylene Rotor and Alkyl Spokes. RSC Adv., 4, 58624-58630.
11. Shinozaki, Y.; Yoshikawa, I.; Araki, K.; Ohara, K.; Yamaguchi, K.; Kawano, S.; Tanaka, K.; Araki, Y.; Wada, T.; *Otsuki, J. (2014). Coordination Oligomers and Polymers of an Oxazole-appended Zinc Chlorophyll

2014

1. Kondo, Y.; *Sasaki, M.; Kawahata, M.; Yamaguchi, K.; Takeda, K. (2014). Enantioselective Synthesis of α -Silylamines by



- Derivative. Chem. Lett., 43, 862-864.
- *Miyamoto, K.; Yokota, Y.; Suefuji, T.; Yamaguchi, K.; Ozawa, T.; Ochiai, M. (2014). Reactivity of Hydroxy- and Aquo(hydroxyl)- λ^3 -iodane-Crown Ether Complexes. Chem. Eur. J., 20(18), 5447-5453.
 - *Setaka, W.; Higa, S.; Yamaguchi, K. (2014). Ring-closing Metathesis for the Synthesis of a Molecular Gyrotop. Org. Biomol. Chem., 12, 3354-3357.
 - *Katagiri, K.; Sakai, T.; Hishikawa, M.; Masu, H.; Masu, H.; Tominaga, M.; Yamaguchi, K.; *Azumaya, I. (2014). Synthesis, Structure, and Thermal Stability of Silver(I) Coordination Polymers with Bis(pyridyl) Ligands Linked by an Aromatic Sulfonamide: One-Dimensional-Straight Chain, One-Dimensional-Columnar with Helical Components, and Two-Dimensional-Layer Network Structures. Cryst. Growth Des., 14, 199-206.
- 2013**
- *Sasaki, M.; Fujiwara, M.; Kotomori, Y.; Kawahata, M.; Yamaguchi, K.; *Takeda, K. (2013). Chirality transfer in Brook rearrangement-mediated S_E2' solvolytic protonation and its use in estimation of the propensity for racemization of the α -lithiocarbanions of the substituents. Tetrahedron, 69, 5823-5828.
 - *Tominaga, M.; Iekushi, A.; Katagiri, K.; Ohara, K.; *Yamaguchi, K. (2013). Organic Crystals Bearing Both Channels and Cavities Formed from Tripodal Adamantane Molecules. Journal of Molecular Structure, 1046, 52-56.
 - *Setaka, W.; Yamaguchi, K. (2013). Order-Disorder Transition of Dipolar Rotor in a Crystalline Molecular Gyrotop and Its Optical Change. J. Am. Chem. Soc., 135, 14560-14563.
 - *Setaka, W.; Koyama, A.; Yamaguchi, K. (2013). Cage Size Effects on the Rotation of Molecular Gyrotops with 1,4-Naphthalenediyl Rotor in Solution. Org. Lett., 15(19), 5092-5095.
 - *Sasaki, M.; Takegawa, T.; Sakamoto, K.; Kotomori, Y.; Otani, Y.; Ohwada, M.; Kawahata, M.; Yamaguchi, K.; Takeda, K. (2013). Enantiodivergent Deprotonation-Acylation of α -Amino Nitriles. Angew. Chem. Int. Ed., 52, 1-6.
 - *Danjo, H.; Iwaso, K.; Kawahata, M.; Ohara, K.; Miyazawa, T.; Yamaguchi, K. (2013). Preparation of Tris(spiroorthocarbonate) Cyclophanes as Back to Back Ditopic Hosts. Org. Lett., 15(9), 2164-2167.
 - *Ohara, K.; Tominaga, M.; Azumaya, I.; *Yamaguchi, K. (2013). Solvent-dependent Assembly of Discrete and Continuous CoCl_2 Adamantane-based Ligand Complexes: Observations by CSI-Mass Spectrometry. and X-ray Crystallography. Anal. Sci., 29(8), 773-776.
 - *Yamaguchi, K. (2013). Cold-Spray Ionization Mass Spectrometry: Applications in Structural Coordination Chemistry. Mass Spec. Soc. J., 2, S0012
 - Shinozaki, Y.; Richard G.; Ogawa, K.; Yamano, A.; Ohara, K.; Yamaguchi, K.; Kawano S.; Tanaka, K.; Araki, Y.; Wada, T.; *Otsuki, J. (2013). Double Helices of a Pyridine-Appended Zinc Chlorophyll Derivative. J. Am. Chem. Soc., 135, 5262-5265.
 - Kawakami, S.; Toyoda, H.; Harinantenaina, L.; Matsunami, *K.; Otsuka, H.; Shinzato, T.; Takeda, Y.; Kawahata, M.; Yamaguchi, K. (2013). Eight New Diterpenoids and Two Nor-Diterpenoids from the Stems of Croton. cascarilloides. Chem. Pharm. Bull., 61(4), 411-418.
 - *Ohwada, T.; Tani, N.; Sakamaki, Y.; Kabasawa, Y.; Otani, Y.; Kawahata, M.; *Yamaguchi, K. (2013). Stereochemical Evidence for Stabilization of a Nitrogen Cation by Neighboring Chlorine or Bromine. Proc. Natl. Acad. Sci. USA, 110, 4206-4211.
 - Uemura, Y.; Sugimoto, S.; Matsunami, K.; Otsuka, H.; *Takeda, Y.; Kawahata, M.; Yamaguchi, K. (2013). Microtropins A-I: 6'-O-(2"S,3"R)-2"-Ethyl-2",3"-dihydroxybutyrates of Aliphatic Alcohol β -D-glucopyranosides from the Branches of Microtropis Japonica. Phytochemistry, 87, 140-147.
 - Masu, H.; Tominaga, M.; Azumaya, I. (2013). Hydrogen-Bonded 1D Chains Formed from Adamantane-Based Bisphenols and Bispyridines: Influences of Substitution Groups on Phenol Ring. Cryst. Growth Des., 13, 752-758.
 - Sakai, T.; Katagiri, K.; Uemura, Y.; Masu, H.; Tominaga, M.; Azumaya, I. (2013). Pseudopolymorphism and Polymorphic Transition Behavior of N-(4'-Methoxyphenyl)-2-naphthalenesulfonamide. Cryst. Growth Des., 13, 308-314.



Laboratory of Biophysics

Staff

Yasushi Kishimoto, Ph.D., Professor
2001 Ph.D. in Biophysics from The University of Tokyo
2001-2003 Postdoctoral Fellow from Mitsubishi Kagaku Institute of Life Sciences to Kanazawa University
2003 Visiting Fellow, NIH/NIHM
2003-2005 JSPS Fellow, Kanazawa University and Osaka University
2006 Assistant Professor, Tokushima Bunri University
2013 Associate Professor, Tokushima Bunri University

Shoji Ueki, Ph.D., Lecturer

Takashi Kubota, Assistant Professor
2005 Ph.D. in Pharmacology from Kyushu University
2005 Assistant Professor, Tokushima Bunri University

Research

Molecular and neural mechanisms of eyeblink classical conditioning

Introduction: Associative learning is a fundamental form of cognition in humans and animals. Eyeblink classical conditioning (EBCC) is a form of associative learning that has been most extensively studied at the neurological and behavioral level. Its basic neural circuitry and neural mechanisms have been demonstrated to be similar in all mammals. Since the same paradigm is applicable to humans as well as non-human mammals, there is growing interest in EBCC for the study of human diseases of motor and memory impairment, in parallel with detailed studies of the molecular and neural mechanisms in animal models.

Typical EBCC experiments use a tone as the conditioned stimulus (CS), and a periorbital shock or corneal air puff as the unconditioned stimulus (US). By repeated presentations of the CS paired with the US, the CS comes to elicit an eyeblink, which is called the conditioned response (CR). Previous studies have indicated that the cerebellum and brainstem are sufficient for *delay* conditioning in which the US is delayed and co-terminates with the CS. On the other hand, *trace* conditioning, in which a stimulus-free trace interval intervenes between the CS and US, requires other brain regions, including the hippocampus and the medial prefrontal cortex (mPFC).

Basic Functions of the Cerebellum and Brainstem: To

investigate the basic properties of the essential neural circuits in the cerebellum and brainstem, we have developed a decerebrate guinea pig preparation, in which a section is made between the thalamus and the superior colliculus and all of the brain tissue above the section is aspirated. Decerebrate animals readily acquire the CR in delay conditioning. When a longer tone CS is used, the learning becomes slower. These CRs are adaptive and appropriately timed relative to the US. Subsequent CS-alone trials cause extinction of the CR. These characteristics of eyeblink conditioning are similar to those reported previously in intact animals of various species, suggesting that the cerebellum and brainstem are sufficient for this type of learning.

We also study trace eyeblink conditioning in decerebrate guinea pigs. A 350-ms tone CS is paired with a 100-ms periorbital shock US with a trace interval of either 0, 100, 250, or 500 ms. Decerebrate animals readily acquire the CR with a trace interval of 0 or 100 ms. Even in the paradigm with a 500-ms trace interval, which is known to depend critically on the hippocampus in all animal species examined, the decerebrate guinea pigs acquire the CR, with the adaptive timing seen in the other paradigms with a shorter trace interval. However, it takes many more trials to learn when we employ the 500-ms trace paradigm rather than the shorter trace-interval paradigms, and the CR expression is unstable from trial to trial. When decerebrate animals are conditioned step by step with a trace interval of 100, 250, and 500 ms (in that order), they easily acquire the adaptive CR with the 500-ms trace interval. However, the CR% decreases after the trace interval is shifted from 250 ms to 500 ms, a decrease that is not observed with the shift from 100 ms to 250 ms. These results suggest that the cerebellum and brainstem can maintain the “trace” of the CS and associate it with the US even in the 500-ms trace paradigm, but that the forebrain might be required to facilitate the association and stabilization of the memory.

Cerebellar Cortical Mechanism of EBCC: Long-term depression (LTD) at parallel fiber-Purkinje cell synapses in the cerebellar cortex has been proposed as the neural substrate for EBCC. Since the glutamate receptor subunit $\delta 2$ (GluR $\delta 2$) is selectively expressed at the dendritic spines of the Purkinje Cell (PC) and is essential for the induction of cerebellar LTD, GluR $\delta 2$ -null mice (in which cerebellar LTD is specifically impaired) provide a useful means to test the cerebellar LTD hypothesis. Mutant mice lacking GluR $\delta 2$ show severe learning impairment in *delay* conditioning, but learn normally

in *trace* conditioning. This surprising finding has now been confirmed in experiments with another line of mutant mice lacking phospholipase C β 4 and with wild-type mice subjected to intracerebellar injection of the NO synthase inhibitor L-NAME, both of which lack cerebellar LTD. Therefore, there may be variations in the cerebellar neural substrates for eyeblink conditioning, depending on the CS-US temporal overlap.

We have recently found that the muscarinic acetylcholine receptor antagonist scopolamine and the NMDA receptor antagonist MK-801 impair learning in trace conditioning experiments with a zero trace-interval (trace 0 paradigm experiments) in GluR δ 2^{-/-} mice, and that the metabotropic glutamate receptor subtype 1 outside the cerebellum is essential for trace conditioning but not for delay conditioning. These findings suggest a contribution of the hippocampus to the LTD-independent learning mechanism. To examine this possibility further, we have looked at the effects of hippocampal lesions on learning in GluR δ 2^{-/-} mice. GluR δ 2^{-/-} mice whose dorsal hippocampi were aspirated exhibit severe learning impairment in the trace 0 paradigm experiments, while control GluR δ 2^{-/-} mice that received a lesion in the cortex overlying the hippocampus are able to learn promptly. Wild-type mice do not show such hippocampal dependency in the trace 0 paradigm. We therefore concluded that the hippocampus is essential for learning with a trace 0 paradigm when cerebellar LTD is disrupted. In contrast, GluR δ 2^{-/-} mice that receive post-training hippocampal lesions retain the memory as well as the control GluR δ 2^{-/-} mice do. The hippocampal lesion also does not affect memory retention in wild-type mice. These results suggest that the hippocampus is *not* essential for *retention* of motor memory with a trace 0 paradigm in LTD-deficient mice. Thus, the present studies clearly argue that the hippocampus is essential for memory formation in cerebellar motor learning when cerebellar LTD is disrupted. These studies suggest that the neural network which underlies learning and memory is both flexible and robust.

Reorganization of Brain Circuitry during Memory

Consolidation: Previous studies, including those described above, have confirmed the time-limited involvement of the hippocampus in mnemonic processes and have suggested that there is reorganization of the responsible brain circuitry during memory consolidation. For temporal characterization of such reorganization, we carried out EBCC experiments with a trace interval of 500 ms in rats with ablation of one of three brain regions: the hippocampus, the mPFC, or the cerebellum. At various time intervals after establishing the trace conditioned response (CR), rats receive an aspiration of one of these regions. After recovery, the animals are tested for their retention of the CR. When ablated one day after the learning, both the hippocampal lesion and the cerebellar lesion groups

exhibit severe impairment in their retention of the CR, whereas the mPFC lesion group show only a small decline. As we increase the interval between the lesion and the learning, the effect of the hippocampal lesion decreases and that of the mPFC lesion increases. When ablated 4 weeks after the learning, the hippocampal lesion group exhibits CRs that are as robust as those of the corresponding control group. In contrast, the mPFC lesion and cerebellar lesion groups fail to retain the CRs. These results indicate that the hippocampus and the cerebellum, but only marginally the mPFC, constitute a brain circuitry that mediates recently acquired memory. As time elapses, the circuitry is reorganized to use mainly the mPFC and the cerebellum, but not the hippocampus, for remotely acquired memory.

Application of EBCC to Studies of Human Memory Loss: In

addition to its great advantage as a model system for learning and memory, EBCC is expected to have wide potential application to clinical studies, including motor impairment (e.g. Parkinson's disease), dementia (e.g. Alzheimer's disease), and other psychopathologies. Such studies using mouse models of human memory loss (experiments using aged animals, senescence-accelerated SAMP8 mice, obese mice and animals injected with β -amyloid peptide) are currently under way.

The Study of the structure and the dynamics of proteins by means of site-directed spin-labeling ESR

The spin label reagent attached to cysteine residue tells us its environment and the structural information of the protein through the ESR spectrum. In other words, the spin label acts as a reporter. In the case of two spin labels introduced into a protein, we can get the distance between two labels by the spectrum. The merit of this method is that we can monitor only the spin label whatever the sample condition is. So we can measure the membrane protein in lipid for example, that is difficult to measure by other spectroscopic method.

Publications

2017

1. Steven A. Connor, Ina Ammendrup-Johnsen, Yasushi Kishimoto, Parisa Karimi Tari, Vedrana Cvetkovska, Takashi Harada, Daiki Ojima, Tohru Yamamoto, Yu Tian Wang, Ann Marie Craig (2017) Altered cortical dynamics and cognitive function upon haploinsufficiency of the autism-linked excitatory synaptic suppressor MDGA2. Cell Rep. 26, 3637-3645.
2. Yasushi Kishimoto, Kai Fukumoto, Mika Nagai, Ayaka Mizuguchi, Yuiko Kobayashi. (2017) Early contextual fear memory deficits in a double-transgenic amyloid- β precursor protein/presenilin 2 mouse model of Alzheimer's disease. Int. J. Alzheimers. Dis. 2017:8584205.

2016

1. Steven A. Connor*, Ina Ammendrup-Johnsen*, Allen W. Chan*,



- Yasushi Kishimoto*, Chiaki Murayama, Naokazu Kurihara, Atsushi Tada, Yuan Ge, Hong Lu, Ryan Yan, Jeffrey M. LeDue, Hirotaka Matsumoto, Hiroshi Kiyonari, Yutaka Kirino, Fumio Matsuzaki, Toshiharu Suzuki, Timothy H. Murphy, Yu Tian Wang, Tohru Yamamoto, Ann Marie Craig. (2016) Altered Cortical Dynamics and Cognitive Function upon Haploinsufficiency of the Autism-Linked Excitatory Synaptic Suppressor MDGA2. *Neuron*, 91, 1052-1068, (2016) (* equal 1st author).
- Hajime Shishido*, Yasushi Kishimoto*, Nobuyuki Kawai, Yasunori Toyota, Masaki Ueno, Takashi Kubota, Yutaka Kirino, Takashi Tamiya. (2016) Traumatic brain injury accelerates amyloid- β deposition and impairs spatial learning in the triple-transgenic mouse model of Alzheimer's disease. *Neurosci. Lett.* 629, 62-67, (* equal 1st author).
 - Yasushi Kishimoto, Hajime Shishido, Mayumi Sawanishi, Yasunori Toyota, Masaki Ueno, Takashi Kubota, Yutaka Kirino, Takashi Tamiya, Nobuyuki Kawai. (2016) Data on amyloid precursor protein accumulation, spontaneous physical activity, and motor learning after traumatic brain injury in the triple-transgenic mouse model of Alzheimer's disease. *Data in Brief* 9, 62-67.
 - Takashi Kubota, Hiroshi Matsumoto, Yutaka Kirino. (2016) Ameliorative effect of membrane-associated estrogen receptor G protein coupled receptor 30 activation on object recognition memory in mouse models of Alzheimer's. *J. Pharmacol. Sci.*, 131, 219-222.
 - Yasushi Kishimoto*, Chiaki Murayama, Naokazu Kurihara, Atsushi Tada, Yuan Ge, Hong Lu, Ryan Yan, Jeffrey M. LeDue, Hirotaka Matsumoto, Hiroshi Kiyonari, Yutaka Kirino, Fumio Matsuzaki, Toshiharu Suzuki, Timothy H. Murphy, Yu Tian Wang, Tohru Yamamoto, Ann Marie Craig. (2016) Altered Cortical Dynamics and Cognitive Function upon Haploinsufficiency of the Autism-Linked Excitatory Synaptic Suppressor MDGA2. *Neuron*, 91, 1052-1068, (2016) (*Equal contribution)
 - Kishimoto Y, Higashihara E, Fukuta A, Nagao A, Kirino Y (2013) Early impairment in a water-finding test in a longitudinal study of the Tg2576 mouse model of Alzheimer's disease. *Brain Res.* 1491:117-126
 - Ishihara Y, Itoh K, Mitsuda Y, Shimada T, Kubota T, Kato C, Song SY, Kobayashi Y, Mori-Yasumoto K, Sekita S, Kirino Y, Yamazaki T, Shimamoto N (2013) Involvement of brain oxidation in the cognitive impairment in a triple transgenic mouse model of Alzheimer's disease: noninvasive measurement of the brain redox state by magnetic resonance imaging. *Free Radic Res.* 47: 731-9
 - Tanimoto, E., Karasawa, S., Ueki, S., Nitta, N., Aoki, I., and Koga, N. (2013) Unexpectedly large water-proton relaxivity of TEMPO incorporated into micelle-oligonucleotides. *RSC Adv* 3, 3531-3534.

2015

- Kishimoto Y, Cagniard B, Yamazaki M, Nakayama J, Sakimura K, Kirino Y, Kano M (2015) Task-specific enhancement of hippocampus-dependent learning in mice deficient in monoacylglycerol lipase, the major hydrolyzing enzyme of the endocannabinoid 2-arachidonoylglycerol. *Front. Behav. Neurosci.* 9:134.
- Kishimoto Y, Yamamoto S, Suzuki K, Toyoda H, Kano M, Tsukada H, Kirino Y (2015) Implicit memory in monkeys: Development of a delay eyeblink conditioning system with parallel electromyographic and high-speed video measurements. *PLoS one* 10:e0129828.

2014

- Ohtani Y, Miyata M, Hashimoto K, Tabata T, Kishimoto Y, Fukaya M, Kase D, Kassai H, Nakao K, Hirata T, Watanabe M, Kano M, Aiba A. (2014) The synaptic targeting of mGluR1 by its carboxyl-terminal domain is crucial for cerebellar function. *J. Neurosci.* 34, 2702-2712.
- Yasuda, S., Yanagi, T., Yamada, M., Ueki, S., Maruta, S., Inoue, A. and Arata, T. (2014) Nucleotide-dependent displacement and dynamics of the alpha-1 helix in kinesin revealed by site-directed spin labeling EPR. *Biophys Biochem Res Commun* 443, 911-916.

2013

- Kishimoto Y, Hirono M, Atarashi R, Sakaguchi S, Yoshioka T, Katamine S, Kirino Y (2013) Age-dependent impairment of eyeblink conditioning in prion protein-deficient mice. *PLOS ONE* 8:e60627
- Kishimoto Y, Kirino Y (2013) Presenilin 2 mutation accelerates the onset of impairment in trace eyeblink conditioning in a mouse model of Alzheimer's disease overexpressing human mutant amyloid precursor protein. *Neurosci. Lett.* 538:15-19.
- Hiasa M*, Isoda Y*, Kishimoto Y*, Saitoh K, Kimura Y, Kanai M, Shibasaki M, Hatakeyama D, Kirino Y, Kuzuhara T (2013) Inhibition of MAO-A and stimulation of behavioural activities



Laboratory of Pharmacognosy and Natural Products Chemistry

Staff

Osamu Shiota, Ph.D.

Professor since 2013

Associate Professor since 2004

Former career: Senior Research Scientist, National Institute of Health Sciences

Ph. D., Graduate School of Pharmaceutical Sciences, Tokyo College of Pharmacy, 1994

Kanami Mori-Yasumoto, Ph. D.

Assistant Professor since 2008

Ph. D., Graduate School of Pharmaceutical Sciences, University of Tokushima, 2008

Research

We conduct chemical and biological research on the components of medicinal plants and crude drugs in order to advance the development of therapeutic pharmaceuticals. The lab focuses on determining the structure of potential compounds, the relationship between that structure and their biological activity, and the role of genes in the biosynthesis of compounds and their function in human physiology. In addition, new avenues in systems biology and metabolomics are being explored and there is ongoing research on the prevention of illegal drug circulation and use, and on the evaluation of drug quality.

I. Research on the biological active chemical constituents of health foods

We found that royal jelly acted at the early stages of the G₁ phase and the S phase of a cell cycle and controlled multiplication of human osteosarcoma cell line, MG-63 cell. Separation of the active water extract by a dialysis membrane and a solid phase extraction suggested that active substances were high polar low molecular compounds. Furthermore, the existence of nitrogen-containing compounds having acidic groups was suggested by LC/MS (ESI) analyses. Further isolation procedure identified that the main active component was AMP N₁-oxide. Continuous examination revealed also the existence of adenosine N₁-oxide, ADP N₁-oxide, ATP N₁-oxide, and NAD N₁-oxide as active ingredients. AMP N₁-oxide and adenosine N₁-oxide inhibited multiplication of MG-63 cell strongly, and their control of the G₁ to S phase comparing with AMP as 1/100 low concentration was found. From these facts, AMP N₁-oxide, adenosine N₁-oxide, and other N₁-oxide are considered to be the main ingredients that contribute at MG-63 cell-growth control of royal jelly.

II. The search of the Alzheimer therapeutic drug from natural resources

Donepezil hydrochloride of the choline esterase inhibitor is used as Dementia and Alzheimer therapeutic drug. Galantamine, the metabolite of the Amaryllidaceae plant acts in similar mechanism, and approval in the country is examined. In addition, Kampo medicine nominates an effect for condition improvement. We perform construction pro-screening to search for the therapeutic drug from a crude drug and a medical plant.

III. The study of anti-Leishmaniasis therapeutics

Leishmaniasis is a parasitic disease caused by species of the genus *Leishmania*. Over 20 of which are known to be pathogenic to humans, and the disease is endemic in some tropical and subtropical regions of the world. *Leishmaniasis* is transmitted by small biting sandflies (*Phlebotomus* spp.), causing a disease which currently afflicts twelve million people in 88 countries. *Leishmania major*, the causative agent of cutaneous leishmaniasis, is a digenetic parasite that exists as an extracellular promastigote within the insect vector (sandfly), and as a nonmotile intracellular amastigote within the phagolysosome of macrophages and other cells of the reticuloendothelial system of the mammalian host. Treatment options for leishmaniasis include pentavalent antimonials as first-line drugs, and amphotericin B and pentamidine as second-line drugs. However, these drugs are extremely toxic and usually too expensive for general use, and more economical and less toxic drugs are thus being sought. We have been searching for plant compounds that are active against *Leishmania major*, *L. panamensis*, *L. guyanensis*, and *L. peruviana*, exhibited significant activity against the pathogenic protozoan, and newly assay method.

IV. Research on the chemical components of illegal drugs of plant origin

1. Khat is a fresh leaf of evergreen shrub *Catha edulis* (Celastraceae) that grows naturally or is grown in Ethiopia, East and Southern Africa, and Yemen, etc., and a lot of people in Africa and Arabia nations use this leaf traditionally as a stimulant biting, and, as a result, it is assumed to obtain the feeling of well-being at the same time as hungry and tiredness's softening. The stimulating component of Khat was believed to be *d*-norpseudoephedrine until cathinone was identified as a main active constituent at the end of 1970's. This cathinone is regulated as narcotics and psychotropic drug, and there is an action similar to (+)-amphetamine that is the synthetic central nervous system stimulation medicine and the strength is assumed to be this level. I synthesized cathinone and ephedrine as an authentic sample to use for the analysis of the drug.

2. *Salvia divinorum* which belongs to Labiatae family is used in

traditional spiritual and curative practices by the indigenous Mazatec people of southern Mexico. Salvinorin A (Sal A), which is the neoclerodane ditrepene and is an extremely potent and highly selective kappa opioid receptor agonist, is the main active constituent isolated from the leaves of *S. divinorum*. The sale of *S. divinorum* has become prohibited due to its psychoactive effect in Japan in recent years. The main objectives of this research are to develop immunoassays using anti-Sal A monoclonal antibody (MAb). The icELISA, which has a measuring range from 0.156 µg/ml to 1.25 µg/ml for Sal A, was developed to distinguish *S. divinorum* among various Labiatae plants. In addition, we are preparing the immunochromatographic strip to realize much more rapid analysis. These immunoassays using anti-Sal A must be a convenient authentication method for *S. divinorum* samples.

V. Research on Development of Preparative Separation Method of Biologically Active Natural Products by Centrifugal Partition Chromatography : Preparative separation of lancemaside A from *Codonopsis lanceolata* by CPC

The roots of *Codonopsis* sp. (Campanulaceae) have been used in folk medicine in China, Korea, and Japan for the treatment of bronchitis, cough, spasm, and inflammation. Recently, it was demonstrated that a hot water extract of *C. lanceolata* roots promoted spermatogenesis and improved sexual motion. Moreover, three phenylpropanoids were identified as the active compounds that promoted spermatogenesis, while several saponins, including lancemaside A, were isolated, and lancemaside A was identified as the active compound that improves sexual motion. Although it is assumed that *C. lanceolata* roots are highly safe since they have been used for a long time, general and specific toxicity tests for safety assurance of the active integrants are required. In general, a large amount of purified compounds is required to assess the effectiveness and to perform safety tests. Therefore we attempted to develop a simple and efficient method for the preparative isolation of lancemaside A from the hot water extract of *C. lanceolata* roots, and resulted in the successful preparative separation of lancemaside A along with two other saponins from the saponin fraction of *C. lanceolata* by CPC.

Publications (2012.4~2017.3)

[Original papers]

2017

1. Tokumura, T., Yoshida, N., Mori-Yasumoto, K., Shirota, O., Kurita, T. (2017). Degradation rates and products of fluticasone propionate in alkaline solutions. *Journal of Pharmaceutical Analysis*, 7(5), 297-302.
2. Kazuma, K., Ando, K., Nihei, K., Wang, X., Rangel, M., Franzolin, M. R., Mori-Yasumoto, K., Sekita, S., Kadowaki, M., Satake, M., Konno, K. (2017). Peptidomic analysis of the venom of the solitary bee *Xylocopa appendiculata circumvolans*, *Journal of Venomous Animals and Toxins including Tropical Diseases*, 23:40.
3. Paudel, M.K., Shirota, O., Sakamoto, S., Morimoto, S., and Tanaka, H. (2017). An immunochromatographic assay for

rapid detection of salvinorin A. *Journal of Immunoassay and Immunochemistry* 38, 438-448.

4. Nugroho, A.E., Okuda, M., Yamamoto, Y., Chin-Piow, W., Hirasawa, Y., Kaneda, T., Shirota, O., Hadi, A.H.A., and Morita, H. (2017). Apowalsogynes A and B, Two Highly Oxidized 3,4-Seco-Apotirucallane Triterpenoids from *Walsura chrysoygne*. *Natural Product Communications* 12, 1189-1192.

2016

1. Muhit, M. A., Umehara, K., Mori-Yasumoto, K., and Noguchi, H. (2016). Furofuran Lignan Glucosides with Estrogen-Inhibitory Properties from the Bangladeshi Medicinal Plant *Terminalia citrina*. *Journal of Natural Products* 79, 1298-1307.
2. Ishihara, Y., Fujitani, N., Sakurai, H., Takemoto, T., Ikeda-Ishihara, N., Mori-Yasumoto, K., Nehira, T., Ishida, A., and Yamazaki, T. (2016). Effects of sex steroid hormones and their metabolites on neuronal injury caused by oxygen-glucose deprivation/reoxygenation in organotypic hippocampal slice cultures. *Steroids* 113, 71-77.
3. Nugroho, A.E., Chin-Piow, W., Hirasawa, Y., Janar, J., Kaneda, T., Shirota, O., and Morita, H. (2016). Daphnane Diterpenoids from *Daphne altaica*. *Natural Product Communications* 11, 1073-1075.

2016

4. Muhit, M.A., Umehara, K., Mori-Yasumoto, K., and Noguchi, H. (2016). Furofuran Lignan Glucosides with Estrogen-Inhibitory Properties from the Bangladeshi Medicinal Plant *Terminalia citrina*. *Journal of Natural Products* 79, 1298-1307.
5. Ishihara, Y., Fujitani, N., Sakurai, H., Takemoto, T., Ikeda-Ishihara, N., Mori-Yasumoto, K., Nehira, T., Ishida, A., and Yamazaki, T. (2016). Effects of sex steroid hormones and their metabolites on neuronal injury caused by oxygen-glucose deprivation/reoxygenation in organotypic hippocampal slice cultures. *Steroids* 113, 71-77.

2015

1. Morishita, Y., Saito, E., Takemura, E., Fujikawa, R., Yamamoto, R., Kuroyanagi, M., Shirota, O., and Muto, N. (2015). Flavonoid glucuronides isolated from spinach inhibit IgE-mediated degranulation in basophilic leukemia RBL-2H3 cells and passive cutaneous anaphylaxis reaction in mice. *Integrative Molecular Medicine* 2, 99-105.
2. Komoto, N., Nakane, T., Matsumoto, S., Hashimoto, S., Shirota, O., Sekita, S., Kuroyanagi, M. (2015). Acyl flavonoids, biflavones, and flavonoids from *Cephalotaxus harringtonia* var. nana. *J Nat Med* 69(4): 479-486.
3. Jenis, J., Nugroho A. E., Hashimoto, A., Deguchi, J., Hirasawa, Y., Chin-Piow, W., Kaneda, T., Shirota, O., Morita, H. (2015). A new benzyloquinoline alkaloid from *Leontice altaica*. *Nat Prod Commun* 10(2): 291-292, 2015.
4. Tsukiura, H., Yamamoto, R., Morishita, Y., Shirota, O., Kuroyanagi, M., Muto, N. (2015). One step formation of capsaicin α -glucoside by enzymatic transglucosylation and its molecular properties. *Japanese Journal of Food Chemistry and Safety*, 22(2), 100-1007.

2014

1. Sallam, A., Nugroho, A.E., Hirasawa, Y., Chin-Piow, W., Kaneda, T., Shirota, O., Gedara, S.R., and Morita, H. (2014). Diterpenoids from *Fagonia mollis*. *Natural Product Communications* 9, 1243-1244.
2. Morita, H., Nugroho, A.E., Nagakura, Y., Hirasawa, Y., Yoshida, H., Kaneda, T., Shirota, O., and Ismail, I.S. (2014). Chrotacumines G-J, chromone alkaloids from *Dysoxylum acutangulum* with osteoclast differentiation inhibitory activity. *Bioorganic & Medicinal Chemistry Letters* 24, 2437-2439.



3. Mori, R., Nugroho, A.E., Hirasawa, Y., Wong, C.P., Kaneda, T., Shirota, O., Hadi, A.H.A., and Morita, H. (2014). Opaciniols A-C, new terpenoids from *Garcinia opaca*. *Journal of Natural Medicines* 68, 186-191.
 4. Kuroyanagi, M., Shirota, O., and Sekita, S. (2014). Transannular cyclization of (4S,5S)-germacrone-4,5-epoxide under basic conditions to yield eudesmane-type sesquiterpenes. *Chemical & Pharmaceutical Bulletin* 62, 725-728.
 5. Kiren, Y., Nugroho, A.E., Hirasawa, Y., Shirota, O., Bekenova, M., Narbekovich, N.O., Shapilova, M., Maeno, H., and Morita, H. (2014). Mumiic acids A-E: new diterpenoids from mumiyo. *Journal of Natural Medicines* 68, 199-205.
 6. Deguchi, J., Sasaki, T., Hirasawa, Y., Kaneda, T., Kusumawati, I., Shirota, O., and Morita, H. (2014). Two novel tetracycles, cassibiphenols A and B from the flowers of *Cassia siamea*. *Tetrahedron Letters* 55, 1362-1365.
 7. Yasumoto, K.; Yasumoto-Hirose, M.; Yasumoto, J.; Murata, R.; Sato, S.; Baba, M.; Mori-Yasumoto, K.; Jimbo, M.; Oshima, Y.; Kusumi, T.; Watabe, S. (2014) Biogenic polyamines capture CO₂ and accelerate extracellular bacterial CaCO₃ formation. *Marine Biotechnology*, 16(4), 465-474.
 8. Morikawa, M., Kino, K., Asada, E., Katagiri, K., Mori-Yasumoto, K., Suzuki, M., Kobayashi, T., Miyazawa, H. (2014). N'-[2-(7,8-Dimethyl-2,4-dioxo-3,4-dihydrobenzo[g]pteridin-10(2H)-yl)ethylidene]-4-nitrobenzohydrazide. *Molbank*, M836; doi: 10.3390/M836.
- 2013**
9. Yamasaki, F., Machida, S., Nakata, A., Nugroho, A.E., Hirasawa, Y., Kaneda, T., Shirota, O., Hagane, N., Sugizaki, T., and Morita, H. (2013). Haworforbins A-C, new phenolics from *Haworthia cymbiformis*. *Journal of Natural Medicines* 67, 212-216.
 10. Paudel, M.K., Shirota, O., Sasaki-Tabata, K., Tanaka, H., Sekita, S., and Morimoto, S. (2013). Development of an Enzyme Immunoassay Using a Monoclonal Antibody against the Psychoactive Diterpenoid Salvinorin A. *Journal of Natural Products* 76, 1654-1660.
 11. Nugroho, A.E., Okuda, M., Yamamoto, Y., Hirasawa, Y., Wong, C.-P., Kaneda, T., Shirota, O., Hadi, A.H.A., and Morita, H. (2013). Walsogynes B-G, limonoids from *Walsura chrysogyne*. *Tetrahedron* 69, 4139-4145.
 12. Hirasawa, Y., Arai, H., Rahman, A., Kusumawati, I., Zaini, N.C., Shirota, O., and Morita, H. (2013). Voacalginines A-E, new indole alkaloids from *Voacanga grandifolia*. *Tetrahedron* 69, 10869-10875.
 13. Fuchino, H., Kiuchi, F., Yamanaka, A., Obu, A., Wada, H., Mori-Yasumoto, K., Kawahara, N., Flores, D., Palacios, O., Sekita, S., Satake, M. (2013). New leishmanicidal stilbenes from a Peruvian folk medicine, *Lonchocarpus nicou*, *Chem Pharm Bull* 61(9), 979-982.
 14. Ishihara, Y., Itoh, K., Mitsuda, Y., Shimada, T., Kubota, T., Kato, C., Song, S., Kobayashi, Y., Mori-Yasumoto, K., Sekita, S., Kirino, Y., Yamazaki, T., Shimamoto, N. (2013). Involvement of brain oxidation in the cognitive impairment in a triple transgenic mouse model of Alzheimer's disease: Non-invasive measurement of the brain redox state by magnetic resonance imaging. *Free Radical Research* 47(9), 731-739.
- 2012**
1. Kuroyanagi, M.; Murata, M.; Nakane, T.; Shirota, O.; Sekita, S.; Fuchino, H.; Shinwari, Z. K. (2012). Leishmanicidal active withanolides from a Pakistani medicinal plant, *Withania coagulans*.: *Chem Pharm Bull* 60(7): 892-897.
 2. Kuroyanagi, M.; Shirota, O.; Sekita, S.; Nakane, T. (2012). Transannular cyclization of (4S,5S)-germacrone-4,5-epoxide into guaiane and secoguaiane-type sesquiterpenes.: *Nat Prod Comm* 7(4): 441-446.
 3. Nugroho, A. E.; Hirasawa, Y.; Piow, W. C.; Kaneda, T.; Hadi, A. H. A.; Shirota, O.; Ekasari, W.; Widyawaruyanti, A.; Morita, H. (2012). Antiplasmodial indole alkaloids from *Leuconotis griffithii*.: *J Nat Med* 66(2): 350-353.
 4. Deguchi, J.; Hirahara, T.; Hirasawa, Y.; Ekasari, W.; Widyawaruyanti, A.; Shirota, O.; Shiro, M.; Morita, H. (2012). New tricyclic alkaloids, cassiarins G, H, J, and K from leaves of *Cassia siamea*.: *Chem Pharm Bull* 60(2): 219-222.
 5. He, F.; Nugroho, A. E.; Wong, C. P.; Hirasawa, Y.; Shirota, O.; Morita, H.; Aisa, H. A. (2012). Rupestines F-M, new guaipyridine sesquiterpene alkaloids from *Artemisia rupestris*.: *Chem Pharm Bull* 60(2): 213-218.
 6. Janar, J.; Nugroho, A. E.; Wong, C. P.; Hirasawa, Y.; Kaneda, T.; Shirota, O.; Morita, H. (2012). Sabiperones A-F, new diterpenoids from *Juniperus Sabina*.: *Chem Pharm Bull* 60(1): 154-159.
 7. Mori-Yasumoto, K., Izumoto, R., Fuchino, H., Ooi, T., Agatsuma, Y., Satake, M., Sekita, S. (2012). Leishmanicidal activities and cytotoxicities of bisnaphthoquinone analogues and naphthol derivatives from Burman *Diospyros burmanica*, *Bioorganic and Medicinal Chemistry* 20, 5215-5219.



Laboratory of Medicinal Chemistry

Staff

Toshie Fujishima, Ph.D.
Professor since 2016
Associate Professor since 2004

Masayuki Morikawa, Ph.D.
Assistant Professor since 2016

Research

Design and synthesis of the novel vitamin D receptor ligands

We have been interested in functions of the nuclear receptors modulated by small molecules, which can be critical to certain disease states. In particular, novel analogues targeted to vitamin D receptor (VDR) were designed and synthesized to understand how the subtype-free, singular VDR can deliver the diverse biological activities vitamin D, as well as to allow the development of potential therapeutic agents with selective activity profiles for the treatment of cancer or osteoporosis. Syntheses of the analogues were carried out by a convergent method using a palladium catalyst. Separate preparation of the requisite A-ring enyne precursors has developed from the 3-buten-1-ol derivatives. Modification in the A-ring, as well as in the side chain of vitamin D, resulted in exceptionally potent compounds with unique activity profiles.

Publications (2013~2017)

[Original papers]

2018

1. Suenaga, T.; Fujishima, T. (2018). The C4-functionalized 9,10-seco-5,7,10(19)-cholestatriene derivatives: Concise synthesis and characterization of novel vitamin D analogues with a four-membered heterocyclic ether, *Tetrahedron*, **2018**, *74*, 1461-1467.

2014

2. Fujishima, T.; Suenaga, T.; Nozaki, T. (2014). Concise synthesis and characterization of novel *seco*-steroids bearing a spiro-oxetane instead of a metabolically labile C3-hydroxy group, *Tetrahedron Lett.*, **2014**, *55*, 3805-3808.
3. Fujishima, T.; Suenaga, T.; Nozaki, T. (2014). Synthetic strategy and biological activity of A-ring stereoisomers of 1,25-dihydroxyvitamin D₃ and C2-modified analogues, *Current Topics Med. Chem.*, **2014**, *14*, 2446-2453.
4. Liu, C.; Zhao, G.-D.; Mao, X.; Suenaga, T.; Fujishima, T.; Zhang, C.-M.; Liu, Z.-P. (2014). Synthesis and biological evaluation of 1 α ,25-dihydroxyvitamin D₃ analogues with aromatic side chains attached at C-17, *Eur. J. Med. Chem.*, **2014**, *85*, 569-575.

2013

5. Fujishima, T.; Nozaki, T.; Suenaga, T. (2013). Design and synthesis of novel 1,25-dihydroxyvitamin D₃ analogues

having a spiro-oxetane fused at the C2 position in the A-ring", *Bioorg. Med. Chem.*, **2013**, *21*, 5209-5217.

[Review]

2016

6. Suenaga, T.; Nozaki, T.; Fujishima, T. (2016). Synthesis of novel vitamin D₃ analogues having a spiro-oxetane structure, *Vitamin*, **2016**, *90*, 109-114.
7. Kino K., Hirao-Suzuki M., Morikawa M., Sakaga A., Miyazawa H. (2017). Generation, repair and replication of guanine oxidation products, *Genes Environ.*, *39*, 21-27.

Laboratory for Molecular Biology

Staff

Hiroshi Miyazawa, Ph. D.

Professor since 2004

Ph.D., The University of Tokyo, 1986

Previous position: Division of Cellular and Gene Therapy Products,
National Institute of Health Sciences, Section Head

Katsuhito Kino, Ph. D.

Associate Professor since 2012

Instructive Professor (Lecturer) since 2004

M.S in Engineering, Graduate School of Engineering, Kyoto
University, 1998

Ph.D. in Medical Science, Tokyo Medical and Dental University,
2002

Previous position: Cellular Physiology Laboratory, RIKEN, Special
postdoctoral researcher at RIKEN

Takanobu Kobayashi, Ph. D.

Assistant Professor since 2015

Ph.D. in Pharmaceutical Sciences, Tokyo University of Science,
2015

Previous position: Graduate School of Pharmaceutical Sciences,
Tokyo University of Science, graduate student

Research

I. Gene expression analysis of mouse embryonic carcinoma P19 cells induced to form neural cells. (Takanobu Kobayashi and Hiroshi Miyazawa)

Mouse embryonic carcinoma P19 cells are pluripotent cells that can be induced to differentiate into multiple cell types by cellular aggregation in the presence of differentiating agents. When aggregated in the presence of all-*trans* retinoic acid (ATRA), P19 cells differentiate into neural cells (including neurons and glia cells), whereas the same cells aggregated in the presence of DMSO differentiate into cardiomyocytes. These cells can simulate the molecular and morphological events occurring during early embryonic development, and have been used extensively as a model to study the molecular mechanisms controlling the process of differentiation into cardiomyocytes or neural cells.

To identify the genes associated with induction of neural differentiation, we carried out a transcriptome analysis of P19 cells induced to form neural cells by ATRA. We employed the DNA microarray method, which can produce an accurate and detailed profile of gene expression. Numerous genes were activated in P19 cells in response to ATRA treatment. We compared the expression profiles from control (undifferentiated) and ATRA-treated P19

cells, which provided an abundance of information on the products involved in neural differentiation. We analyzed the sequential expression patterns of some genes during neural differentiation. We are investigating the relationship between ATRA treatment and expression patterns of these genes, interactions with other factors, and function in neural differentiation. These findings will provide useful information for a comprehensive understanding of the complex process of the induction of neural differentiation. We are conducting additional investigations to better understand the role of these genes during the induction of neural differentiation.

Recently, we identified a candidate gene associated with the process of neural differentiation. We found that the expression of this gene is induced by all-*trans* retinoic acid (ATRA) during neural differentiation in P19 cells from our study using microarray analysis. We describe the induction mechanism of this gene and its activation in this process. In conclusion, the expression of this gene is upregulated via ATRA-bound RAR α and binding to the RARE in the Csn3 promoter region. This will be a first step forward in unraveling the mysteries of the role of this gene in the process of neural differentiation.

II. DNA Oxidation, Point Mutation and DNA Damage (Kino)

The genome is constantly assaulted by oxidative stress, and DNA lesions are likely to be associated with oxygen metabolism. DNA lesions are generated by many of these oxidative stressors. Genotoxic alterations in the genomic DNA are implicated in aging and in several pathophysiological processes, particularly those associated with cancer.

1. Guanine Oxidation

Photosensitized oxidation of guanine produces reactive oxygen species, including 8-oxoguanine, which is a mutagenic DNA lesion. Riboflavin (vitamin B2) is known as a photosensitizer for the oxidation of guanine. We have demonstrated: the user-friendly synthesis and purification of a flavin-linked oligonucleotide; the practical synthesis of hydroxyethyl-flavin from commercially available riboflavin; the preparation of a flavin-linked oligonucleotide containing phosphoramidite of hydroxyethyl-flavin. To evaluate the usefulness of this method, the flavin-linked oligonucleotide was synthesized. The flavin-linked oligonucleotide and oligonucleotide containing 8-oxoG were then irradiated (366 nm) at neutral pH. Enzymatic digestion of the irradiated mixture indicated that the 8-oxoG residue



Regulation of Immune Functions through Nuclear Receptors

Staff

Professor: Yoshiharu Ohoka, Ph.D.

Associate Professor since 2005

Ph.D. Tokyo Institute of Technology, 1993

Research Associate: Aya Yokota-Nakatsuma, Ph.D.

Assistant Professor since 2006

Ph.D. Kitasato University, 2004

Research

Research Themes:

The mechanisms of immune cell trafficking and the regulation of immune responses are our main themes to clarify. Especially, we study roles of nuclear receptor ligands including vitamin A & D and various hormones in regulating immune functions especially in mucosal systems including the gut. By pursuing these biologically fundamental questions, we set a goal to establish a solid basis of new remedies and drug discovery for various diseases.

Recent Study:

For efficient immune responses, immune cells with proper functions need to migrate into right sites in the body. T cells, known as the control tower of the immune system, patrol the whole body along with the blood vessels and lymphatic vessels. However, they cannot migrate into non-lymphoid tissues before they are activated with antigen in the secondary lymphoid organs. Once they are activated and become effector or memory T cells, however, they can migrate into non-lymphoid tissues. They tend to migrate into the tissue that is associated with the secondary lymphoid organ where they are activated. This type of migration is called “homing”. For example, T cells that are activated with antigen in the small intestine-related secondary lymphoid organs, Peyer’s patches (PP) and mesenteric lymph nodes (MLN), tend to migrate into small intestinal tissues including the lamina propria. In 2004, we found that vitamin A-derived retinoic acid is the physiological factor that imprints gut-homing specificity on T cells. We also found that subpopulations of dendritic cells (DC) in PP and MLN express the key enzyme of retinoic acid synthesis, RALDH (retinaldehyde dehydrogenase), and are capable of producing retinoic acid from vitamin A (retinol). They imprint T cells with the gut-homing specificity by delivering retinoic acid to T cells during antigen presentation. In 2006, we also found that a similar mechanism is involved in the imprinting of B cells with gut-homing specificity by a collaboration mainly with Dr. von Andrian’s group and Dr. Adams’ group.

In 2009, we established a method for estimating the enzyme activity of RALDH in each DC, and identified the retinoic acid-producing subpopulation in MLN-DC and PP-DC. The RALDH2 isoform was mostly responsible for the activity. Depending on these results, we searched for the physiological factors that induce RALDH2 expression in DC in the gut or in MLN. We found that GM-CSF (granulocyte-macrophage colony-stimulating factor) plays a major role in the induction, and that retinoic acid itself plays a role as an essential cofactor. IL-4 and IL-13 exhibited effects similar to those of GM-CSF on RALDH2 expression, but are found to be dispensable by the analysis of their receptor-deficient mice. The stimulation through Toll-like receptors enhanced the RALDH2 expression in DC as well as DC maturation.

In 2007, several groups reported that the retinoic acid-producing DC also enhance the differentiation of Foxp3⁺ inducible regulatory T cells (iTreg) and suppress that of pro-inflammatory Th17 cells. Accordingly, we found that GM-CSF-treated DC that expressed RALDH2 could enhance the differentiation of Foxp3⁺ iTreg and suppress that of Th17. These results suggest that retinoic acid contributes to oral tolerance and regulation of immune responses to specific antigens. We have recently found that vitamin A deficiency affects not only the nature of T cells but also that of DC, and that MLN-DC in vitamin A deficient mice can induce oral antigen-specific CD4⁺ T cells that produce high levels of IL-13 and tumor necrosis factor- α (TNF- α). Under vitamin A-deficient conditions, although it has been generally known that antibody responses are reduced, we found that markedly high levels of IgG1 antibody responses and IgE antibody responses against oral antigens can be induced. It is likely that these antibody responses involve the above-mentioned new IL-13-producing inflammatory helper T (Th) cells. Currently, we are investigating the molecular mechanism of differentiation of these Th cells and their role in allergic and inflammatory diseases.

We have been also analyzing some other aspects of retinoic acid effects on immune functions and regulation, including 1) The molecular mechanism of retinoic acid effects on the expression of gut-homing receptors on immune cells, 2) The role of a retinoic acid-metabolizing system in the regulation of T cell functions, 3) Amplification and disruption of retinoic acid signals by RXR ligands and environmental chemicals, 4) The molecular mechanism of RALDH isoform 2 (RALDH2, encoded by *Aldh1a2*) expression in DC and the roles of a retinoic acid-bound RAR/RXR heterodimer and a retinoic acid response element (RARE) half-site at the proximal promoter of the *Aldh1a2* gene. The RARE half-site

in this gene promoter was commonly found in many animal species.

We found that, in vitamin A-deficient mice, dextran sulfate sodium (DSS) induced more severe colitis, and a higher rate of development of colorectal carcinoma with colitis following treatment with azoxymethane, compared with vitamin A-supplemented mice. Therefore, vitamin A is likely to inhibit the development of chronic inflammation and cancer. These vitamin A effects may also be dependent on retinoic acid production. As we mentioned above, GM-CSF and retinoic acid itself play important roles in the development of retinoic acid-producing DC, and Toll-like receptor-mediated stimulation enhances their maturation and retinoic acid-producing capacity. However, we recently found that Toll-like receptor stimulation also induced production of inflammatory cytokines. As retinoic acid-producing RALDH2^{high} MLN-DC in steady-state mice do not produce inflammatory cytokines, we searched for a stimulation condition that can induce maturation in GM-CSF/retinoic acid-treated semi-mature DC and can enhance their RALDH2 expression without inducing inflammatory cytokines. We found that stimulation with immobilized proteins such as E-cadherin/IgG-Fc chimeric protein fulfilled the requirement. These proteins induced signals through integrin β 1 in DC. E-cadherin expressed on epithelial cells in gut tissues might contribute to the development of the RALDH2^{high} DC. These DC induced in vitro could efficiently induced gut-homing iTreg, and could significantly suppress DSS-induced colitis.

Publications

* 2013-2017

2016

1. Yokota-Nakatsuma, A., Ohoka, Y., Takeuchi, H., Song, S.-Y., and Iwata, M.: Beta 1-integrin ligation and TLR ligation enhance GM-CSF-induced ALDH1A2 expression in dendritic cells, but differentially regulate their anti-inflammatory properties. *Sci Rep.* 6:37914 (2016).
2. Okayasu, I., Hana, K., Nemoto, N., Yoshida, T., Saegusa, M., Yokota-Nakatsuma, A., Song, S.-Y., and Iwata, M.: Vitamin A inhibits development of dextran sulfate sodium-induced colitis and colon cancer in a mouse model. *BioMed Res Int*: Article ID 4874809 (2016).

2014

1. Ohoka, Y., Yokota-Nakatsuma, A., Maeda, N., Takeuchi, H., and Iwata, M.: Retinoic acid and GM-CSF coordinately induce retinal dehydrogenase 2 (RALDH2) expression through cooperation between the RAR/RXR complex and Sp1 in dendritic cells. *PLoS One.* 9(5):e96512 (2014).
2. Yokota-Nakatsuma, A., Takeuchi, H., Ohoka, Y., Kato, C., Song, S.-Y., Hoshino, T., Yagita, H., Ohteki, T., and Iwata, M.: Retinoic acid prevents mesenteric lymph node dendritic cells from inducing IL-13-producing inflammatory Th2 cells.

Mucosal Immunol. 7(4):786-801 (2014). Epub 2013 Nov 13.

2013

1. Takeuchi, H., Yokota-Nakatsuma, Y., Ohoka, Y., Kagechika, H., Kato, C., Song, S.-Y., and Iwata, M.: Retinoid X receptor agonists modulate Foxp3⁺ regulatory T cell and Th17 cell differentiation with differential dependence on retinoic acid receptor activation. *J Immunol* 191(7):3725-3733 (2013). Epub 2013 Aug 26.
2. Sato, T., Kitawaki, T., Fujita, H., Iwata, M., Iyoda, T., Inaba, K., Ohteki, T., Hasegawa, S., Kawada, K., Sakai, Y., Ikeuchi, H., Nakase, H., Niwa, A., Takaori-Kondo, A., and Kadowaki, N.: Human CD1c⁺ myeloid dendritic cells acquire a high level of retinoic acid-producing capacity in response to vitamin D₃. *J Immunol* 191(6):3152-3160 (2013). Epub 2013 Aug 21.

[Review articles in Japanese]

2017

1. Yokota-Nakatsuma, A. Retinoic acid prevents dendritic cells from inducing novel inflammatory T cells that produce abundant interleukin-13. *Yakugaku Zasshi* 137(12):1491-1496(2017).

2014

1. Yokota-Nakatsuma, A. and Iwata, M. Regulation of inflammatory dendritic cells by vitamin A. *Inflammation & Immunity* (Ensho-to-Men-eki) 22(4):63(295)-67(299) (2014).

2013

1. Yokota-Nakatsuma, A. Regulation of retinoic acid-producing ability in dendritic cells. *Clinical Immunology & Allergy* (Rinsho Men-eki Allergy-ka) 59(3):392-397 (2013).



Laboratory of Pharmaceutical Health Sciences

Staff

Hiromi Nochi, Ph.D.

Professor since April, 2013.

Associate Professor (April, 2006 - March, 2013)

Previous position: Lecturer at Faculty of Pharmaceutical Sciences,
Health Sciences University of Hokkaido.

Hajime Takeuchi, Ph.D.

Associate Professor since 2013

Previous position: Postdoctrand at University of Zurich
(Switzerland)

Yoshimitsu Kiriyama, Ph.D.

Assistant Professor since 2005.

Previous position: Postdoctoral Researcher at McGill University
Health Centre (Canada).

Research

Research project: Analysis of the molecular mechanism by which extracellular acidification induces inflammatory responses in rheumatoid arthritis synovial cells. (Nochi)

The local acidification of extracellular pH is caused by augmentation of cell proliferation as observed in the cancer tissue and the inflammatory site. In RA, the proliferation of synovial cell is abnormally augmented and the pH of synovial fluid from RA patient is lower than that of normal synovial fluid. Therefore, we examined the possibility that the intraarticular acidification may affect the inflammatory responses and contributes to exacerbation of pathological condition in RA. We found that extracellular acidic pH induced ADAMTS-4 expression through Gq-coupled proton-sensing receptor (OGR1) in RA synovial cells. Furthermore, extracellular acidic pH-induced and ADAMTS-4 expression were markedly inhibited by siRNA targeted for the OGR1 receptor and the inhibitors for G_{q/11} protein and p38 MAPK. We conclude that the OGR1/ G_{q/11}/p38 MAPK or NFκB pathway regulates ADAMTS-4 expression under acidic condition. To examine details of the intracellular signaling mechanism by which ADAMTS-4 induction are regulated under acidic circumstances, further studies are now under way.

Research project: characterization of retinoid X receptor signaling on T cell differentiation. (Takeuchi)

Retinoic acid (RA) is an immune-modulating molecule, and its signaling is known to affect T cell differentiation. It can enhance differentiation toward regulatory T cell (Treg), and suppress that

toward Th17. RA receptor is consist of two components, RAR and RXR. Although RXR has a ligand-binding domain, it does not bind RA at physiological condition. It is not well-studied whether RXR-specific signaling can affect T cell differentiation.

We found that RXR signaling actually played some roles on T cell differentiation. RXR signaling dramatically enhanced RA-mediated Treg differentiation. On the other hand, it suppressed that of Th17 in cooperation with some nuclear receptor signaling. The effect of RXR signal was observed in vivo as well as in vitro. Thus, this finding can apply to develop new methods to regulate immune-response and inflammatory diseases.

Research project: Analysis of the molecular mechanism of GABARAPs in autophagy. (Kiriyama, Nochi)

Astrocytes play several important roles in the central nervous system: providing nutrients to neurons, forming the blood-brain barrier, maintaining the extracellular ion balance, surrounding and maintaining synapses, and participating with neurons in the processing of information. Thus, optimal function of the brain depends on maintenance of homeostasis of astrocytes as well as neurons from stresses. Cell survival and maintenance of homeostasis depends on removal of damaged cellular cytosolic components. Macroautophagy (hereafter referred to as autophagy) is the process by which defective proteins and damaged organelles are isolated and degraded to generate new proteins and organelles under a variety of stresses. Moreover, impairment of autophagic processes results in neurodegeneration

We found that reactive oxygen species (ROS) and the mitochondrial damage leads to the induction of autophagic flux and the upregulation of gamma-aminobutyric-acid-type-A receptor-associated protein (GBARAP) subfamily proteins in C6 glioma cells. GABARAPL1 mRNA among GABARAP subfamily members were highly upregulated by ROS. GABARAPL1 are associated with both autophagosome formation and carrying target proteins to autophagosomes. Thus, the expression of the GABARAP subfamily members may depend on discrete regulations of the expression of mRNA and protein, and the GABARAP subfamily members may have discrete functions in autophagy. Further studies on the mechanism of GABARAP subfamily members in autophagy are under way.

Publications (2012~2017)

[Original papers]

2018

1. Kiriyama Y. and Nochi H. Intra- and Intercellular Quality

Control Mechanisms of Mitochondria. *Cells*.7:1, (2018)

2017

2016

1. Yokota-Nakatsuma, A, Ohoka, Y., Takeuchi, H., Song, S.-Y. and Iwata, M. Beta1-integrin ligation and TLR ligation enhance GM-CSF-induced ALDH1A2 expression in dendritic cells, but differentially regulate their anti-inflammatory properties. *Sci Rep.* 6, 37914 (2016).
2. Kiriyama, Y., Kasai, K., Kino, K. & Nochi, H. Induction of the expression of GABARAPL1 by hydrogen peroxide in C6 glioma cells. *Integr Mol Med* 3, 675-679 (2016).
3. Kiriyama, Y. & Nochi, H. D-Amino Acids in the Nervous and Endocrine Systems. *Scientifica (Cairo)*, 6494621 (2016).

2015

1. Kiriyama, Y., Kino, K., and Nochi, H. Autophagy and amino acids with their metabolites. *Integr Food Nutr Metab* 2:151-155 (2015).
2. Kiriyama, Y., Ozaki, A., Kino, K., and Nochi, H. Effects of CCCP on the expression of GABARAPL2 in C6 glioma cells. *Integr Mol Med* 2: 177-180 (2015).
3. Kino, K., Nakatsuma, A., Nochi, H., Kiriyama, Y., Kurita, T., Kobayashi, T., and Miyazawa, H. Commentary on the Phototoxicity and Absorption of Vitamin B2 and Its Degradation Product, Lumichrome. *Pharm Anal Acta* 6:403 (2015).
4. Kiriyama, Y., and Nochi, H. The Function of Autophagy in Neurodegenerative Diseases. *Int J Mol Sci* 16: 26797-26812 (2015).
5. Nakatsuma, A., Wada, S., Kamano, J., Kiriyama, Y., Kino, K., and Ninomiya, M. The effects of herbal teas on drug permeability. *Integr Mol Med* 3: 453-456 (2015).
6. Nakatsuma, A., Kiriyama, Y., Kino, K., and Ninomiya, M. Diabetes drugs that protect pancreatic β cells. *Integr Mol Med* 3: 467-472 (2015).

2014

1. Ohoka, Y, Yokota-Nakatsuma, A, Maeda, N, Takeuchi, H. and Iwata, M.: Retinoic acid and GM-CSF coordinately induce retinal dehydrogenase 2 (RALDH2) expression through cooperation between the RAR/RXR complex and Sp1 in dendritic cells. *PLoS One.* 9(5): e96512 (2014).
2. Yokota-Nakatsuma, A., Takeuchi, H., Ohoka, Y., Kato, C., Song, S.-Y., Hoshino, T., Yagita, H., Ohteki, T., and Iwata, M.: Retinoic acid prevents mesenteric lymph node dendritic cells from inducing IL-13-producing inflammatory Th2 cells. *Mucosal Immunol.* 7(4):786-801 (2014). Epub 2013 Nov 13.

2013

1. Takeuchi, H., Yokota-Nakatsuma, Y., Ohoka, Y., Kagechika, H., Kato, C., Song, S.-Y., and Iwata, M.: Retinoid X receptor agonists modulate Foxp3⁺ regulatory T cell and Th17 cell differentiation with differential dependence on retinoic acid receptor activation. *J Immunol* 191(7): 3725-3733 (2013).



Staff

Maki K. Yamada, Ph. D.

Professor since 2016

Ph. D. in The University of Tokyo, 1996

Shuntaro Kohnomi Ph. D.

Assistant Professor since 2012, till Jun 2017

Ph. D. in Okayama University, 2009

Kazuya Kuboyama Ph. D.

Assistant Professor since June 2017

Ph. D. in Kyushu University, 2011

Research

Main Project (Maki K. Yamada, Kazuya Kuboyama, Shuntaro Kohnomi)

Seeing through the brain mechanism for memory using transgenic mice

Background

We showed that an F-actin stabilizing protein CapZ accumulated and stayed around synapses (spines) that underwent long-term potentiation (LTP) (Genes to Cells 2010), thus the place for memory-related change in neuronal synapses is expected to be labeled by EGFP-CapZ even in-vivo. On the other hand, we published another paper (Cerebral cortex 2009) suggesting that memory-coding neurons are included in a part of Arc-expressing neurons, thus, Yamada made the transgenic mouse of EGFP-CapZ driven by the Arc-promotor, Arc::EGFP-CapZ, to mark the memory coding synapses and neurons. In the resulting mouse having blight fluorescence and normal learning ability, the EGFP-fluorescence is mainly found in some of synaptic structure, spines, and the cell body.

Till now,

1. In-vitro confirmation; It was confirmed that the LTP-inducing stimuli increased the level of the green fluorescence in a part of the synaptic structure (spines) and the neuronal cell body in the hippocampus sections from the TG having Arc::EGFP-CapZ (ACE-TG) as expected.

2. In-vivo observation; we observed that a small number of strongly green neurons emerged after the visually-cued learning in the primary visual cortex of the ACE-TG.

3. Ex-vivo analyses; we analyzed the changes of the green neurons and spines in the primary visual cortex by making fixed brain sections from ACE-TG with or without the same visual learning and then found the differences.

Future plans: Seeing through the brain mechanism for memory by multidisciplinary approach.

References from Yamada's group

1. Cereb. Cortex

Experience-dependent, rapid structural changes in hippocampal pyramidal cell spines.

Kitanishi T, Ikegaya Y, Matsuki N, Yamada MK*

Cerebral cortex (New York, N.Y.) 19(11) 2572-2578 (2009)

<http://cercor.oxfordjournals.org/content/19/11/2572.short>

2. Genes Cells

Activity-dependent localization in spines of the F-actin capping protein CapZ screened in a rat model of dementia.

Kitanishi T, Sakai J, Kojima S, Saitoh Y, Inokuchi K, Fukaya M,

Watanabe M, Matsuki N, Yamada MK* Genes to cells : devoted

to molecular & cellular mechanisms 15(7) 737-747 (2010)

<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2443.2010.01411.x/pdf>

Publications (2012~2017)

[Original papers]

2017

1. Kuboyama K, Tanga N, Suzuki R, Fujikawa A, Noda M. (2017) Protamine neutralizes chondroitin sulfate proteoglycan-mediated inhibition of oligodendrocyte differentiation. *PLoS one* 12: e0189164.
2. Fujikawa A, Chow JPH, Matsumoto M, Suzuki R, Kuboyama K, Yamamoto N, Noda M. (2017) Identification of novel splicing variants of protein tyrosine phosphatase receptor type Z. *J biochem* 162: 381-390.
3. Fujikawa A, Sugawara H, Tanaka T, Matsumoto M, Kuboyama K, Suzuki R, Tanga N, Ogata A, Masumura M, Noda M. (2017) Targeting PTPRZ inhibits stem cell-like properties and tumorigenicity in glioblastoma cells. *Sci rep* 7: 5609.
4. Kohnomi S, Ebihara K, Kobayashi M. (2017) Suppressive regulation of lateral inhibition between medium spiny neurons via dopamine D₁ receptors in the rat nucleus accumbens shell. *Neurosci Lett* 636: 58-63.
5. Nakahata Y, Eto K., Murakoshi H, Watanabe M, Kuriu T, Hirata H, Moorhouse AJ, Ishibashi H, Nabekura J. (2017) Activation-dependent rapid postsynaptic clustering of glycine receptors in mature spinal cord neurons. *eNeuro* 4: e0194-16.2017.

2016

6. Kuboyama K, Fujikawa A, Suzuki R, Tanga N, Noda M. (2016) Role of Chondroitin Sulfate (CS) Modification in the Regulation of Protein-tyrosine Phosphatase Receptor Type Z (PTPRZ) Activity. PLEIOTROPHIN-PTPRZ-A SIGNALING IS INVOLVED IN OLIGODENDROCYTE DIFFERENTIATION. *J Biol Chem* 291: 18117-18128.
7. Fujikawa A, Nagahira A, Sugawara H, Ishii K, Imajo S, Matsumoto M, Kuboyama K, Suzuki R, Tanga N, Noda M, Uchiyama S, Tomoo T, Ogata A, Masumura M, Noda M. (2016) Small-molecule inhibition of PTPRZ reduces tumor growth in a rat model of glioblastoma. *Sci rep* 6: 20473.
8. Ono Y, Saitow F & Konishi S. (2016) Differential modulation of GABA_A receptors underlies postsynaptic depolarization- and

Laboratory of Pharmaceutical Education

purinoceptor-mediated enhancement of cerebellar inhibitory transmission: a nonstationary fluctuation analysis study. *PLoS one* 11: e0150636.

2015

9. Ikeda, T., Makino, Y., and Yamada, M.K. (2015) 17alpha-Estradiol Is Generated Locally in the Male Rat Brain and Can Regulate GAD65 Expression and Anxiety, *Neuropharmacology*, Vol. 90.p.9-14
10. Hamamoto, H., Urai, M., Ishii, K., Yasukawa, J., Paudel, A., Murai, M., Kaji, T., Kuranaga, T., Hamase, K., Katsu, T., Su, J., Adachi, T., Uchida, R., Tomoda, H., Yamada, M., Souma, M., Kurihara, H., Inoue, M., and Sekimizu, K. (2015) Lysocin E Is a New Antibiotic That Targets Menaquinone in the Bacterial Membrane. *Nat. Chem. Biol.*, Vol. 11.p.127-33
11. Kuboyama K., Fujikawa A, Suzuki R, Noda M. (2015) Inactivation of protein tyrosine phosphatase receptor type Z by pleiotrophin promotes remyelination through activation of differentiation of oligodendrocyte precursor cells. *J neurosci* 35: 12162-12171.
12. Matsumoto M, Hiyama TY, Kuboyama K., Suzuki R, Fujikawa A, Noda M. (2015) Channel properties of Nax expressed in neurons. *PLoS one* 10: e0126109.
13. Fujikawa A, Matsumoto M, Kuboyama K., Suzuki R, Noda M. (2015) Specific dephosphorylation at tyr-554 of git1 by ptpz promotes its association with paxillin and hic-5. *PLoS one* 10: e0119361.
14. Almuriexhi M, Shintani T, Fahiminiya S, Fujikawa A, Kuboyama K., Takeuchi Y, Nawaz Z, Nadaf J, Kamel H, Kitam AK, Samiha Z, Mahmoud L, Ben-Omran T, Majewski J, Noda M. (2015) Loss-of-function mutation in APC2 causes Sotos syndrome features. *Cell rep* 10: 1585-1598.
15. Kohnomi S., Konishi S. (2015) Multiple actions of a D₃ dopamine receptor agonist, PD128907, on GABAergic inhibitory transmission between medium spiny neurons in mouse nucleus accumbens shell. *Neurosci Lett* 600: 17-21.
16. Shoji M, Arakaki Y, Esumi T, Kohnomi S., Yamamoto C, Suzuki Y, Takahashi E, Konishi S., Kido H, Kuzuhara T. (2015) Bakuchiol is a phenolic isoprenoid with novel enantiomer-selective anti-influenza A virus activity involving Nrf2 activation. *J Biol Chem* 290: 28001-28017.
17. Okabe A, Shimizu-Okabe C, Arata A, Konishi S., Fukuda A, Takayama C. (2015) KCC2-mediated regulation of respiration-related rhythmic activity during postnatal development in mouse medulla oblonga. *Brain Res* 1601: 31-39.

2014

18. Suzuki R, Matsumoto M, Fujikawa A, Kato A, Kuboyama K., Yonehara K, Shintani T, Sakuta H, Noda M. (2014) SPIG1 negatively regulates BDNF maturation. *J Neurosci* 34: 3429-3442.
19. Isshiki M, Tanaka S, Kuriu T., Tabuchi K, Takumi T, Okabe S. (2014) Enhanced synapse remodeling as a common phenotype in mouse models of autism. *Nat Commun* 5: 4742.

2013

20. Shin E, Kashiwagi Y, Kuriu T., Iwasaki H, Tanaka T, Koizumi H, Gleeson JG, Okabe S. (2013) Doublecortin-like kinase enhances dendritic remodeling and negatively regulates synapse maturation. *Nat Commun* 4: 1440.

2012

21. Matsumoto M, Fujikawa A, Suzuki R, Shimizu H, Kuboyama K., Hiyama TY, Hall RA, Noda M. (2012) SAP97 promotes the stability of Nax channels at the plasma membrane. *FEBS Lett* 586: 3805-3812.
22. Kuboyama K., Fujikawa A, Masumura M, Suzuki R,

Matsumoto M, Noda M. (2012) Protein tyrosine phosphatase receptor type z negatively regulates oligodendrocyte differentiation and myelination. *PLoS one* 7: e48797.

23. Hirono M, Saitow F, Kudo M, Suzuki H, Yanagawa Y, Yamada M, Nagao S, Konishi S., Obata K. (2012) Cerebellar Globular Cells Receive Monoaminergic excitation and monosynaptic inhibition from Purkinje cells. *PLoS one* 7: e29663.
24. Kohnomi S., Koshikawa N, Kobayashi M. (2012) D₂-like dopamine receptors differentially regulate unitary IPSCs depending on presynaptic GABAergic neuron subtypes in rat nucleus accumbens shell. *J Neurophysiol* 107: 692-703.
25. Kuriu T., Yanagawa Y, Konishi S. (2012) Activity-dependent coordinated mobility of hippocampal inhibitory synapses visualized with presynaptic and postsynaptic tagged-molecular markers. *Mol Cell Neurosci* 49: 184-195.

[Review article]

26. Yamada MK. (2016) Angiogenesis in refractory depression: A possible phenotypic target to avoid the blood brain barrier. *Drug Discov Ther* 10: 79-81.
27. Yamada, MK. (2016) A link between vascular damage and cognitive deficits after whole-brain radiation therapy for cancer: A clue to other types of dementia? *Drug Discov Ther* 10: 74-78.
28. Konishi S., Satake S. (2013) Physiological interactions between neurons and glia: roles of transporters in the control of intersynaptic crosstalk. Chapter 9, In *Glial Cells: Embryonic Development, Types, Functions and Role in Disease*, edit by K. Charanjit and Eng-Ang Ling, pp.177-191, Nova Science Publishers, New York.

[Books]

29. Konishi S., Kirino Y, Ito E, Song SY. (2013) "Mechanism of Memory" Vol. 1 & 2: translated from "Memory From Mind to Molecules" by E.R. Kandel & L.R. Squire. pp.1-298, pp.1-300, Blubacks B-1842 & B-1843, Kodansha, Tokyo

Patent

30. Fujikawa A, Kuboyama K., Noda M. patent number: JP 6238252 B



Laboratory of Pathological Physiology.

Staff

Si-Young Song, M.D.,
Professor since April 1, 2006
Visiting Scientist of Tokyo Metropolitan Institute for Neuroscience
Doctor of Medical Science of Tokyo Medical and Dental University,
1983

Kentaro Nakashima
Research assistant since Nov. 1, 2006
Master of Engineering of Yokohama National University, 2005

Shintaro Iwashita
Visiting Professor since April 1, 2017
Doctor of Philosophy of University of Tokyo, 1975

Rie Fujii
Laboratory assistant since Nov 1, 2013 until April 30, 2017

Tomoko Warita
Laboratory assistant since April 1, 2017
Doctor of Agriculture of Kobe University,

Research

Research Themes:

It's clear that the future direction of the Pharmaceutical science is "the development of therapeutic method of human diseases based on the understanding of pathophysiology at molecular level". This direction is also one of the most important bases of Pharmaceutical education. Thus we set final goal of our research on the integrative understanding of the pathogenesis of human diseases from molecular to individual level. To pursue this task, animal models for human diseases are beneficial experimental tools. From the analyses of these animals, we have a chance to combine analyses of molecular and cellular level with clinical changes at individual level. Further, these animals are valuable, because we can obtain most early changes in the pathogenesis of diseases, which are hardly examined in human patients. Thus one of the main methods of our division is histopathological analysis of these animals. These analyses require systemic approaches from macroscopic anatomy, conventional histological methods, immuno histochemical and *in situ* hybridization histochemical methods using light microscopy and electron microscopy. Now we are trying to establish a system for analyses integrating these lesions identified by histological techniques with biochemical and molecular biological analyses. We also pursue cooperative research projects with medical institutions outside our university for analyses using human materials that fulfill ethical criteria, depending on the progress in each research project. These trials using good animal models for human disease are expected to contribute to the better understanding of the pathophysiology of

human disease. Following are the detailed information of each research project.

Analysis of functional role of lanosterol 14 α -demethylase (LDM) in remyelination following cuprizone-induced demyelination using transgenic mice that highly express LDM.

Pathogenesis of demyelinating disease such as multiple sclerosis is not yet clear and effective therapy is not established. Lipid is a major component of myelin and remyelination involves various kinds of lipid metabolism- related enzymes, which can be therapeutic targets of demyelinating disease. Cholesterol, a major component of myelin, cannot pass blood brain barrier and all of cholesterol necessary for myelin formation is believed to be synthesized in the brain. Our previous study revealed that the expression of lanosterol 14 α -demethylase (LDM, CYP51) is up-regulated during the process of remyelination. LDM catalyzes demethylation of lanosterol to FF-MAS (Follicular fluid meiosis-activating sterol) in cholesterol biosynthesis and is the only cytochrome P450 enzyme that is involved in cholesterol biosynthesis in eukaryotes. We previously found that LDM is selectively expressed in oligodendroglia. We succeeded to establish a line of transgenic mouse that shows oligodendroglia-specific high expression of LDM (LDM-Tg). We disclosed following points using this transgenic mice and wild type mice.

1. Western blot analysis

Cerebral white matter was collected by Laser Microdissection from unfixed brain of LDM-Tg and wild type mice. Western blot of isolated proteins from the cerebral white matter was done using anti-LDM and anti-proteolipid protein (PLP) antibody.

- (1) The expression of LDM and PLP showed a tendency to decrease at the demyelination stage after treatment with cuprizone for 6 weeks and increase at remyelination stage after changing to normal diet for 1 or 3 weeks in both LDM-Tg and control mice. But the degree of expression change was different between both mice.
- (2) Expression of LDM was remarkably high in LDM-Tg as compared with wild type and its expression changes were mild at demyelination and remyelination stage and generally maintained the level of control treated with normal diet for 6 weeks. Especially decrease in the expression of LDM and PLP was apparently mild as compared with wild type.

2. Immunohistochemistry

Frozen sections were made from fixed brain of LDM-Tg and wild type mice. Immunohistochemical analysis was done using anti-LDM, anti-PLP, anti-Iba-1 and anti-GFAP antibody. Myelin

staining was done using FluoroMyelin Green (FMG).

- (1) Demyelination was significantly mild in LDM-Tg as compared with wild type after treatment with cuprizone for 6 weeks by myelin staining, suggesting that reactive remyelination that occur during the process of demyelination is more accelerated in LDM-Tg than wild type. Difference in the degree of demyelination between both mice was most apparent at demyelination stage and not so apparent after changing to normal diet, where remyelination is rapidly progress also in wild type mice.
- (2) In accordance with the finding of myelin staining, decrease in the PLP-immunoreactivity, increase in the Iba-1-immunoreactivity that reflect response of microglia treating collapsed myelin and the GFAP-immunoreactivity that reflect response of astrocytes were mild in LDM-Tg as compared with wild type. Difference in the immunoreactivity between both mice was also most apparent at demyelination stage was most apparent at demyelination stage and not so apparent after changing to normal diet.

These two kinds of experiment suggest that overexpression of LDM accelerates remyelination through increase in cholesterol biosynthesis in oligodendroglia during the process of experimental demyelination induced by the treatment with cuprizone. If we can develop drugs that can enhance the expression and/or enzyme activity of LDM, they can be effective therapeutic tools for demyelinating disease

Studies on a common epitope between two non-homologous proteins, Bcnt/Cfdp1 and glutamine synthetase.

We had reported that an antibody against a peptide, EELAIHNRGKEGYIERKA, of the evolutionarily conserved C-terminal domain of Bcnt/Cfdp1 (anti-BCNT-C Ab) specifically cross-reacts with glutamine synthetase (GS, glutamate-ammonia ligase) by Western blotting of brain extracts, while other anti-Bcnt/Cfdp1 antibodies do not. No significant homology of amino acid sequences was found between Bcnt/Cfdp1 and GS. However, we identified a key amino acid sequence, KKGYFE, localized in a limited region of the C-terminal of GS, by a combination of Ab binding analyses using deletion mutants, amino acid-substituted mutants of GS and binding competition using peptide fragments of GS. We considered KEGYIEA of the antigen peptide as its counterpart. Furthermore, substitution of Phe-337 of GS with several amino acids suggests that Phe-337 is not directly engaged in the epitope-paratope interface responsible for the cross-reactivity with the anti-BCNT-C Ab, but its steric bulk is important to maintain the stereochemical environment of the interface. These findings show that the anti-BCNT-C Ab recognizes two distinct molecules, Bcnt/Cfdp1 protein and GS through a common spatial linear epitope and that such a cross-reactivity is mainly due to the physicochemical environment formed by amino acid residues of the epitope rather than the primary sequence of the

epitope, giving a novel insight into the specificity of antigen-antibody reaction and protein-protein interaction.

Publications

2016

1. Okayasu I, Hana K, Nemoto N, Yoshida T, Saegusa M, Yokota-Nakatsuma A, Song S-Y, and Iwata M. Vitamin A inhibits development of dextran sulfate sodium-induced colitis and colon cancer in a mouse model. *BioMed Res Int* Article ID 4874809 (2016). [Epub 2016 May 19].
2. Yokota-Nakatsuma A, Ohoka Y, Takeuchi H, Song S-Y, and Iwata M. Beta 1-integrin ligation and TLR ligation enhance GM-CSF-induced ALDH1A2 expression in dendritic cells, but differentially regulate their anti-inflammatory properties. *Sci Rep* 6:37914. doi: 10.1038/srep37914 (2016)

2015

1. Iwashita S, Suzuki T, Yasuda T, Nakashima K, Sakamoto T, Kohno T, Takahashi I, Kobayashi T, Ohno-Iwashita Y, Imajoh-Ohmi S, Song SY, Dohmae N. Mammalian Bcnt/Cfdp1, a potential epigenetic factor characterized by an acidic stretch in the disordered N-terminal and Ser250 phosphorylation in the conserved C-terminal regions. *Biosci. Rep.* 35: 1-12 (2015)

2014

1. Yokota-Nakatsuma A, Takeuchi H, Ohoka Y, Kato C, Song S-Y, Hoshino T, Ohteki T and Iwata M. Retinoic acid prevents mesenteric lymph node dendritic cells from inducing IL-13-producing inflammatory Th2 cells. *Mucosal Immunol* 7:786-801 (2014).

2013

1. Ishihara Y, Itoh K, Mitsuda Y, Shimada T, Kubota T, Kato C, Song S-Y, Kobayashi Y, Mori-Yasumoto K, Sekita S, Kirino Y, Yamazaki T, Shimamoto N. Involvement of brain oxidation in the cognitive impairment in a triple transgenic mouse model of Alzheimer's disease: noninvasive measurement of the brain redox state by magnetic resonance imaging. *Free Radic Res* 47: 731-739 (2013).
2. Takeuchi H, Yokota-Nakatsuma A, Ohoka Y, Kagechika H, Kato C, Song S-Y and Iwata M. Retinoid X Receptor Agonists Modulate Foxp3 + Regulatory T Cell and Th17 Cell Differentiation with Differential Dependence on Retinoic Acid Receptor Activation. *J Immunol* 191: 3725 -3733 (2013)
3. Yokota-Nakatsuma A, Takeuchi H, Ohoka Y, Kato C, Song S-Y, Hoshino T, Ohteki T and Iwata M. Retinoic acid prevents mesenteric lymph node dendritic cells from inducing IL-13-producing inflammatory Th2 cells. *Mucosal Immunol* Epub 2013 Nov 13.



Laboratory for Pharmacotherapy and Experimental Neurology

Staff

Kouichi Itoh, Ph. D.

Professor since April 01, 2004.

M. Sc., Showa College of Pharmaceutical Sciences graduate school of pharmacology, 1983.

Ph. D., Toho University, Sch. of Med. 1991

Previous occupation: The Tokyo metropolitan organization for medical research, Tokyo Metropolitan Institute of Medical Science, the division of pharmacology, Researcher.

Taira Matsuo, Ph.D.

Lecturer since 2015

Assistant Professor since 2009

Ph.D. Okayama University, 2009

Rie Komori, Ph. D.

Assistant Professor since 2005.

D.Sc. Nara Women's University, 2003.

Previous occupation: Department of Etiology and Pathophysiology, National Cardiovascular Center Research Institute, postdoctoral researcher.

Research

【Research aims】

Our research goal is the novel molecular target for new antiepileptic drugs. To achieve this goal, we are working on molecular mechanism for the epileptogenesis of partial and generalize epilepsy through the several approaches such as pharmacological, behavioral, cell biological, biochemical and *in vivo* imaging techniques.

【Research Scopes】

1. Prevention of status epilepticus-induced brain edema and neuronal cell loss by new antiepileptic drugs.

Status epilepticus (SE) refers to neurologic emergencies that may lead to death or permanent neurologic injury. To avoid life threatening injury, patients must be properly and quickly treated. Furthermore, SE causes 3~5% of symptomatic epilepsy (~35% of epileptic syndromes), thus SE patients are at a high risk of developing acquired epilepsy (Hesdorffer, 1998; Temkin, 2003; Jacobs et al., 2009). The management of SE is important to prevent mortality and the development of post-SE symptomatic epilepsy. Seizures must be treated as soon as possible and

benzodiazepines (lorazepam or diazepam) are typically administered as first-line antiepileptic drugs (AEDs). However, when these drugs fail, second-line AEDs (phenytoin; PHT, fosphenytoin; fosPHT, valproate; VPA, and midazolam) are administered in refractory SE prior to giving phenobarbital; PB (Manno, 2011). Various clinical trials have indicated that conventional AEDs (e.g., DZP, PB, VPA, or PHT) suppressed acute seizures, but thus far there has been no success at preventing the development of post-SE acquired epilepsy under various conditions (Temkin, 2001; 2003; 2009). Although the mechanisms underlying the development of acquired epilepsy as part of the epileptogenic process are not well understood, the lack of efficacy of the AEDs suggests that the biological mechanisms of the acquired epilepsy process may be quite different from that of the established epileptic brain (Pitkanen et al., 2009).

Levetiracetam ([[(S)- α -ethyl-2-oxo-1-pyrrolidine acetamide]]) with broad-spectrum antiepileptic effects is an established second-generation AED that is widely used in patients with either generalized or partial epilepsy (Lyseng-Williamson, 2011). In addition, levetiracetam is one of currently available candidates as second-line AED for SE (Manno, 2011) and as an anti-epileptogenic drug (Pearl et al., 2013; Klein, et al., 2012). Animal studies have shown that levetiracetam possesses anticonvulsant activity and results in neuroprotective effects (Mazarati et al., 2004; Zheng et al., 2010). In addition, levetiracetam has been considered for the treatment of pilocarpine (PILO)-SE due to its anti-epileptogenic effects in basic and clinical studies. Two phase II clinical trials for levetiracetam indicated the possibility that it may decrease the risk of acquired epilepsy or prevent the development of acquired epilepsy (Pearl et al., 2013; Klein, et al., 2012). However, the previous evidence in SE animal models has been conflicting and whether levetiracetam can prevent or modify epileptogenesis remains controversial (Löscher et al., 1998; Glien 2002; Klitgaard, Pitkanen, 2003; Stratton et al., 2003; Gibbs et al., 2006; Brandt, et al., 2007).

Temporal lobe epilepsy (TLE) is the most frequent type (75%) of symptomatic partial epilepsies that originate from the limbic regions (e.g., hippocampus and amygdala) after an initial brain insult, such as SE, stroke, and traumatic brain injury (TBI). Additionally, it is also one of the most refractory forms of epilepsy with approximately 30% of patients being unresponsive to AEDs (Engel, 1996; Kwan and Brodie, 2000). In this present study, we used PILO-induced SE mice as a model of TLE to determine the effects of repeated administration of high-dose levetiracetam after the termination of SE by DZP. We observed that repeated

high-dose levetiracetam prevented the development of brain edema in the limbic regions at the initial period of post-SE, and the incidence of spontaneous recurrent seizures. In the present study, we determined the possible molecular and cellular mechanisms of LEV treatment after termination of SE. To assess the effect of LEV against the brain alterations after SE, we focused on blood-brain barrier (BBB) dysfunction associated with angiogenesis and brain inflammation. The consecutive treatment of LEV inhibited the temporarily increased BBB leakage in the hippocampus two days after SE. At the same time point, the LEV treatment significantly inhibited the increase in the number of CD31-positive endothelial immature cells and in the expression of angiogenic factors. These findings suggested that the increase in neovascularization led to an increase in BBB permeability by SE-induced BBB failure, and these brain alterations were prevented by LEV treatment. Furthermore, in the acute phase of the latent period, pro-inflammatory responses for epileptogenic targets in microglia and astrocytes of the hippocampus activated, and these upregulations of pro-inflammatory-related molecules were inhibited by LEV treatment. These findings suggest that LEV is likely involved in neuroprotection via anti-angiogenesis and anti-inflammatory activities against BBB dysfunction in the acute phase of epileptogenesis after SE.

2. Study on the relationship between blood brain barrier (BBB) disruption and epilepsy.

In recent years, it has been well recognized that the therapies for epilepsy by the AEDs, which is represented by valproate, have been definitely effective. On the other hand, no less than 30% of epileptic patients were intractable, so there are difficulties in achievement of high level of QOL for them. In order to dissolve this problem, the development of new AEDs with novel mechanisms is an important for drug-resistant patients. We aim to find out the novel molecular target for new drugs. Recently, we have focused the relationship between BBB disruption and generalized epilepsy. Although conventional evaluation methods of BBB disruption are to measure the diffusion of low molecular weight dye (ex. Evans blue) to brain parenchyma, they are not available in animals alive. In our laboratory, the spatial and sequential changes of the BBB disruption in PTZ-induced convulsive alive mice were elucidated by the technique of gadolinium-enhanced magnetic resonance imaging using the MRI for rodent (MRminiSR, 1.5T). In addition, we have investigated the involvement of NO in the BBB disruption in generalized epilepsy.

Publications (2013~2018)

[Original papers]

2018

1. M. Tanaka, Y. Ishihara, S. Mizuno, A. Ishida, C. F. Vogel, M.

Tsuji, T. Yamazaki, **K. Itoh**. (2018) Progression of vasogenic edema induced by activated microglia under permanent middle cerebral artery occlusion. *Biochemical and Biophysical Research Communications*, 496:582-587.

2. Kotani, M., Sato, Y., Ueno, A., Ito, T., **Itoh, K.**, Imadae, M. (2018) Two non-cytotoxic type 2 ribosome-inactivating proteins (Sambucus Sieboldiana lectin and Sambucus Nigra lectin) lead neurosphere cells to caspase-independent apoptosis. *Biomedical Research* accepted

2017

3. Yasuhiro Ishihara, **Kouichi Itoh**, Miki Tanaka, Mayumi Tsuji, Toshihiro Kawamoto, Suguru Kawato, Christoph F. A. Vogel, Takeshi Yamazaki. (2017) Potentiation of 17 β -estradiol synthesis in the brain and elongation of seizure latency through dietary supplementation with docosahexaenoic acid. *Scientific Reports* 7: 6268 | DOI:10.1038/s41598-017-06630-0.

2016

4. Kotani, M., Sato, Y., Ueno, A., Ito, T., **Itoh, K.**, Imadae, M. (2016) A Novel Monoclonal Antibody against Neuroepithelial and Ependymal Cells and Characteristics of Its Positive Cells in Neurospheres. *Cell Mol Neurobiol*. 36:11-26.
5. **Itoh, K.**, Ishihara, Y., Komori, R., Nochi, H., Taniguchi, R., Chiba, Y., Ueno, M., Takata-Tsuji, F., Dohgu, S., Kataoka, Y. (2016) Levetiracetam treatment influences blood-brain barrier failure associated with angiogenesis and inflammatory responses in the acute phase of epileptogenesis in post-status epilepticus mice. *Brain Res.*, 1652:1-13

6. **Kamada M.**, Mitsui Y., **Matsuo T.**, Takahashi T. (2016) Reversible transformation and de-differentiation of human cells derived from induced pluripotent stem cell teratomas. *Hum Cell*. 29(1):1-9

2015

7. Ishihara Y., **Itoh K.**, Ishida A., Yamazaki T. (2015) Selective estrogen-receptor modulators suppress microglial activation and neuronal cell death via an estrogen receptor-dependent pathway. *J. Steroid Biochem. Mol. Biol.* 145:85-93.
8. **Itoh, K.**, Inamine, M., Oshima, W., Kotani, M., Chib, Y., Ueno, M., Ishihara, Y., (2015) Prevention of status epilepticus-induced brain edema and neuronal cell loss by repeated treatment with high-dose levetiracetam. *Brain Res.* 1608:225-234. Correspondence
9. **Itoh, K.**, Mizuno, S., Hama, S., Oshima, W., Kawamata, M., Hossain, A., Ishihara, Y., Tokuda, M. (2015) Beneficial effects of supplementation of the rare sugar "D-allulose" against hepatic steatosis and severe obesity in Lepob/Lepob mice. *J. Food Sci.* 80: H1619-H1626. Correspondence



10. Ishihara, Y., Takemoto, T., **Itoh, K.**, Ishida, A., Yamazaki, T. (2015) Dual Role of Superoxide Dismutase 2 Induced in Activated Microglia: Oxidative Stress Tolerance and Convergence of Inflammatory Responses *J. Biol. Chem.* 290:22805-22817.

(2013) Csn3 gene is regulated by all-trans retinoic acid during neural differentiation in mouse P19 cells. *Plos One* 8, e61938.

2014

11. Kobayashi T., **Komori R.**, Ishida K., Kino K., Tanuma S., Miyazawa H., (2014) Tal2 expression is induced by all-trans retinoic acid in P19 cells prior to acquisition of neural fate. *Sci. Rep.*, 4, 4935
12. Kamada M., Mitsui Y., Kumazaki T., Kawahara Y., **Matsuo T.**, Takahashi T. (2014) Tumorigenic risk of human induced pluripotent stem cell explants cultured on mouse SNL76/7 feeder cells. *Biochem Biophys Res Commun.* 453(3):668-73
13. **Matsuo T.**, Ogawa W., Tsuchiya T., Kuroda T. (2014) Overexpression of *vmeTUV* encoding a multidrug efflux transporter of *Vibrio parahaemolyticus* causes bile acid resistance. *Gene.* 541(1):19-25.
14. Kumazaki, T., Takahashi, T., **Matsuo, T.**, Kamada, M., and Mitsui, Y. (2014) Re-emergence of undifferentiated cells from transplants of human induced pluripotent stem cells as a possible risk factor of tumorigenesis. *Cell Biol. Int. Rep.* 21(1):17-24

2013

15. Danjo, S., Ishihara, Y., Watanabe, M., Nakamura, Y., **Itoh, K.** (2013) Pentylentetrazole-induced loss of blood-brain barrier integrity involves excess nitric oxide generation by neuronal nitric oxide synthase. *Brain Res.* 1530, 44-53.
16. Watanabe, M., Miyai, A., Danjo, S., Nakamura, Y., **Itoh, K.** (2013) The threshold of pentylentetrazole-induced convulsive seizures, but not that of nonconvulsive seizures, is controlled by the nitric oxide levels in murine brains. *Exp. Neurol.* 247, 645-652.
17. Fujii H. G., Sato-Akaba H., Emoto M. C., **Itoh K.**, Ishihara Y., Hirata H., (2013) Noninvasive mapping of the redox status in septic mouse by in vivo electron paramagnetic resonance imaging. *Magn. Reson. Imag.* 31:130-138
18. Ishihara Y, **Itoh K.**, Mitsuda Y, Shimada T, Kubota T, Kato C, Song Si-Y, Kobayashi Y, Mori-Yasumoto K, Sekita S, Kirino Y, Yamazaki T, Shimamoto N. (2013) Involvement of brain oxidation in the cognitive impairment in 3xTg-AD mice: Non-invasive measurement of the brain redox state by magnetic resonance imaging. *Free Radical Res.* 47: 731-739
19. **Matsuo T.**, Nakamura K, Kodama T, Mikami T, Hiyoshi H, Tsuchiya T, Ogawa W, Kuroda T. Characterization of all RND-type multidrug efflux transporters in *Vibrio parahaemolyticus*. *Microbiologyopen.* (2013) 2(5):725-742.
20. **Komori R.**, Kobayashi T, Matsuo H, Kino K, Miyazawa H.



Laboratory of Pharmacokinetics and Pharmacodynamics

Staff

Yoshihisa Kato, Ph.D., Pharmacist

Professor since 2010

Visiting Professor of University of Shizuoka

Ph.D. University of Shizuoka, 1991

Norikazu Sakakibara, Ph. D., Pharmacist

Lecturer since 2012

Postdoctoral at Department of Applied Life Science, Kyoto University

Ph.D. Kyoto University, 2005

Kazutaka Atobe, Ph. D., Pharmacist

Assistant Professor since 2007

Ph.D. The University of Tokushima, 2007

Research

1. Integrated study on pharmacokinetics and pharmacodynamics for efficient drug discovery and on the optimum drug therapy
2. Basic and clinical analysis of drug-metabolizing enzyme and transporters influencing pharmacokinetics and pharmacodynamics
3. Basic research on the toxicity mechanism of drugs, xenobiotics and their active metabolites
4. Basic research for synthesis of COA-Cl (2-Cl-C.OXT-A) analogs, and their tube formation activities of human umbilical vein endothelial cells (HUVEC)
5. Design, synthesis, and anti-HBV activity of heterocyclic derivatives
6. The development of novel reaction for the heterocyclic bioactive substances
7. Curcumin and doxorubicin co-encapsulation liposome effects on cell growth, apoptosis, and angiogenesis

The elucidation of relationship between pharmacokinetics (PK) and pharmacodynamics (PD) of drugs is critical not only for the discovery of novel drugs but also for their optimum uses in clinical settings. The effects of clinically used drugs are influenced by drug-metabolizing enzymes and drug-transporters. Therefore, one of our major research projects is to push an integrated analysis of PK and PD for investigational drugs by characterizing *in vivo* drug-metabolizing enzymes and drug-transporters under physiological and pathological conditions.

We are also conducting the research of mechanism(s) for the decrease in levels of serum thyroid hormones by xenobiotics and their active metabolites, and the species difference and

extrapolation to human in these compounds-induced alteration of the hormone levels.

In addition to these projects, we have also focused on simultaneous determination of polybrominated diphenyl ethers (PBDEs), hydroxylated (OH-) and methoxylated (MeO-) PBDEs found in marine sponge by atmosphere pressure chemical ionization tandem mass spectrometry (APCI-LC/MS/MS), and immunotargeting of liposome to tumor and endothelial cell expressed membrane type-1 matrix metalloproteinase (MT1-MMP).

Stimulator of angiogenesis: Stimulators of angiogenesis are sometimes desired for clinical treatment of diseases evoked by an impaired blood supply including ulcers associated with diabetes or burn wounds. However, availability of stimulators is few till date because of their size. Most of the stimulators known are endogenous large molecules like VEGF and FGF. Those are expensive proteins, hard to synthesize, and not so stable. We developed 2-Cl-C.OXT-A as a stable candidate compound. This compound strongly stimulates the tube formation of HUVEC. Its maximum potency at 100 μ M was stronger than VEGF (10ng/mL). 2-Cl-C.OXT-A will be applicable to medicine instead of endogenous growth factors such as VEGF and/or FGF.

Effect of uracil analog against HIV-1: Human immunodeficiency virus type 1 (HIV-1) contains an important enzyme, reverse transcriptase (RT), which catalyzes the conversion of the viral genome RNA into the double-stranding DNA. Since this process is essential for viral replication, many drugs targeting this enzyme have been developed. Within the class of the anti-HIV agents which inhibit reverse transcriptase, non-nucleoside reverse transcriptase inhibitors (NNRTIs) are rapidly increasing. It is interesting that some NNRTIs have an aromatic group at the 6 position of uracil. Under the background of these reports, we undertook a search for an anti-HIV agent by the SAR of the 1,3-disubstituted uracil. Two compounds showed excellent anti-HIV-1 activity with moderate cytotoxicity.

Publications (2013-2018)

[Original papers]

2018

1. Fujii, Y., Kato, Y., Sakamoto, K., Matsuishi, T., Harada, H.K., Koizumi, A., Kimura, O., Endo, T., and Haraguchi, K. (2018). Tissue-specific bioaccumulation of long-chain perfluorinated carboxylic acids and halogenated methylbipyrroles in Dall's porpoises (*Phocoenoides dalli*) and harbor porpoises (*Phocoena phocoena*) stranded in northern Japan. *Sci. Total Environ.*, 616-617, 554-563.
2. Fujii, Y., Kato, Y., Masuda, N., Harada, K., Koizumi, A., and Haraguchi, K. (2018) Contamination trends and factors affecting the transfer of hexabromocyclododecane

diastereomers, tetrabromobisphenol A, and 2,4,6-tribromophenol to breast milk in Japan. *Environ. Pollut. online* 20 March.

2017

1. Kato, Y., Fujii, A., Haraguchi, K., Fujii, Y., Atobe, K., Endo, T., Kimura, O., Koga, N., Ohta, C., Yamada, S., and Degawa, M. (2017). Possible mechanism for the polychlorinated biphenyl induced liver-selective accumulation of thyroxine in rats. *J. Toxicol. Sci.*, 42, 663-669.
2. Endo, T., Kimura, O., Terasaki M., Fujii, Y., Haraguchi, K., Ohta, C., Koga, N., and Kato, Y. (2017). Growth-related changes in non-essential and essential metals in the liver of star-spotted smooth-hounds (dogfish) *Mustelus manazo* from the northern region of Japan. *Mar. Environ. Res.* 131, 156-161.
3. Kimura, O., Fujii, Y., Haraguchi, K., Kato, Y., Ohta, C., Koga, N., and Endo, T. (2017). Uptake of perfluorooctanoic acid by Caco-2 cells: Involvement of organic anion transporting polypeptides. *Toxicol. Lett.*, 277, 18-23.
4. Sakakibara, N., Igarashi, J., Takata, M., Konishi, R., Kato, Y., and Tsukamoto, I. (2017). Synthesis and evaluation of novel cyclopropane nucleoside as potential tube formation agents. *Chem. Pharm. Bull.*, 65, 504-510.
5. Ohta, C., Fujii, Y., Haraguchi, K., Kato, Y., Kimura, O., Endo, T., and Koga, N. (2017). Metabolism of 2,2',3,4,4',5,6'-heptachlorobiphenyl (CB182) by rat, guinea pig and human liver microsomes. *Fukuoka Acta Medica* 108: 51-57.

2016

1. Endo, T., Kimura, O., Ohta, C., Koga, N., Kato, Y., Fujii, Y., and Haraguchi, K. (2016). Metal concentrations in the liver and stable isotope ratios of carbon and nitrogen in the muscle of silvertip shark (*Carcharhinus albimarginatus*) culled off Ishigaki Island, Japan: changes with growth. *PLOS ONE*, 11(2):e0147797. doi: 10.1371/journal.pone.0147797.
2. Kimura, O., Fujii, Y., Haraguchi, K., Ohta, C., Koga, N., Kato, Y., Endo, T. (2016). Effect of quercetin on the uptake and efflux of aristolochic acid I from Caco-2 cell monolayers. *J. Pharm. Pharmacol.*, 68, 883-889.
3. Igarashi, J., Okamoto, R., Yamashita, T., Hashimoto, T., Karita, S., Nakai, K., Kubota, Y., Takata, M., Yamaguchi, F., Tokuda, M., Sakakibara, N., Tsukamoto, I., Konishi, R., Hirano, K. (2016). A key role of PGC-1 α transcriptional coactivator in production of VEGF by a novel angiogenic agent COA-Cl in cultured human fibroblasts. *Physiological reports*, 4, e12742.

2015

1. Ohta, C., Haraguchi, K., Kato, Y., Endo, T., Kimura, O., and Koga, N. (2015). Distribution and excretion of 2,2',3,4',5,5',6'-heptachlorobiphenyl (CB187) and its metabolites in rats and guinea pigs. *Chemosphere*, 118, 5-11.
2. Endo, T., Kimura, O., Ogasawara, H., Ohta, C., Koga, N., Kato, Y., and Haraguchi, K. (2015). Mercury, cadmium, zinc and copper concentrations and stable isotope ratios of carbon and nitrogen in tiger sharks (*Galeocerdo cuvier*) culled off Ishigaki Island, Japan. *Ecol. Indic.*, 55, 86-93.
3. Tanino, T., Funakami, Y., Nagai, N., and Kato, Y. (2015). Cyclosporin A-sensitive cytotoxicity of flurbiprofen non-stereoselectively mediated by cytochrome P450 metabolism in three-dimensional cultured rat hepatocytes. *J Pharm Pharmacol*, 67, 1406-1415.
4. Sakakibara, N., Balboni, G., Congiu, C., Onnis, V., Demizu, Y., Misawa, T., Kurihara, M., Kato, Y., Maruyama, T., Okamoto, M., Toyama, M., and Baba, M. (2015). Design, synthesis, and anti-HIV-1 activity of 1-substituted 3-(3,5-dimethylbenzyl)triazine derivatives. *Antiviral Chemistry & Chemotherapy*, 24, 62-71.
5. Sakakibara, N., Igarashi, J., Takata, M., Konishi, R., Suzue, N., Kato, Y., Maruyama, T., and Tsukamoto, I. (2015). Design, synthesis, and evaluation of novel 2-halogenated or aminated

carbocyclic oxetanocin A analogs as potential angiogenic agents. *Heterocycles*, 91, 1823-1832.

6. Sakakibara, N., Igarashi, J., Takata, M., Demizu, Y., Misawa, T., Kurihara, M., Konishi, R., Kato, Y., Maruyama, T., and Tsukamoto, I. (2015). Synthesis and evaluation of novel carbocyclic oxetanocin A (COA-Cl) derivatives as potential tube formation agents. *Chem. Pharm. Bull.*, 63, 701-709.
7. Sakakibara, N., Baba, M., Okamoto, M., Toyama, M., Demizu, Y., Misawa, T., Kurihara, M., Irie, K., Kato, Y., and Maruyama, T. (2015). Design, synthesis, and anti-HIV-1 activity of 1-aromatic methyl-substituted 3-(3,5-dimethylbenzyl)uracil and N-3,5-dimethylbenzyl-substituted urea derivatives. *Antiviral Chemistry & Chemotherapy*, 24, 3-18.
8. Ohta, C., Haraguchi, K., Kato, Y., Endo, T., Kimura, O., and Koga, N. (2015). Metabolism of 2,2',3,4,4',5,5'-Heptachlorobiphenyl (CB180) by animal liver microsomes. *Fukuoka Acta Medica* 106: 176-183.

2014

1. Kato, Y., Haraguchi, K., Onishi, M., Ikushiro, S., Endo, T., Ohta, C., Koga, N., Yamada, S., and Degawa, M. (2014). 3,3',4,4'-Tetrachlorobiphenyl-mediated decrease of serum thyroxine level in C57BL/6 and DBA/2 mice occurs mainly through enhanced accumulation of thyroxine in the liver. *Biol. Pharm. Bull.*, 37, 504-509.
2. Kimura, O., Ohta, C., Koga, N., Haraguchi, K., Kato, Y., and Endo, T. (2014). Carrier-mediated uptake of nobiletin, a citrus polymethoxyflavonoid, in human intestinal Caco-2 cells. *Food Chemistry*, 154, 145-150.
3. Fujii, Y., Nishimura, E., Kato, Y., Harada, K.H., Koizumi, A., and Haraguchi, K. (2014). Dietary exposure to phenolic and methoxylated organohalogen contaminants in relation to their concentrations in breast milk and serum in Japan. *Environ. Int.*, 63, 19-25.
4. Matsubara, F., Sagara, Y., Kato, Y., Harada, K., Koizumi, A., and Haraguchi, K. (2014). Detection of antibodies to human T-cell leukemia virus types 1 and 2 in breast milk from East Asian women. *Biol. Pharm. Bull.*, 37, 311-314.
5. Kimura, O., Haraguchi, K., Ohta, C., Koga, N., Kato, Y., and Endo, T. (2014). Uptake of aristolochic acid I into Caco-2 cells by the monocarboxylic acid transporters. *Biol. Pharm. Bull.*, 37, 1475-1479.
6. Igarashi, J., Hashimoto, T., Kubota, Y., Shoji, K., Maruyama, T., Sakakibara, N., Takuwa, Y., Ujihara, Y., Katanosaka, Y., Mohri, S., Naruse, K., Yamashita, T., Okamoto, R., Hirano, K., Kasaka, H., Takata, M., Konishi, R., Tsukamoto, I. (2014). Involvement of S1P₁ receptor pathway in angiogenic effects of a novel adenosine-like nucleic acid analog COA-Cl in cultured human vascular endothelial cells. *Pharmacology Research & Perspectives*, 2, e00068.

2013

1. Kato, Y., Onishi, M., Haraguchi, K., Ikushiro, S., Ohta, C., Koga, N., Endo, T., Yamada, S., and Degawa, M. (2013). A possible mechanism for 2,3',4,4',5'-pentachlorobiphenyl-mediated decrease in serum thyroxine level in mice. *Biol. Pharm. Bull.*, 36, 1594-1601. (Highlighted paper selected by Editor-in-Chief)
2. Hidaka, N., Suemaru, K., Kato, Y., and Araki, H. (2013). Involvement of $\alpha 4\beta 2$ nicotinic acetylcholine receptors in working memory impairment induced by repeated electroconvulsive seizures in rats. *Epilepsy Research*, 104, 181-185.
3. Endo, T., Hisamichi, Y., Kimura, O., Ogasawara, H., Ohta, C., Koga, N., Kato, Y., and Haraguchi, K. (2013). Levels of mercury in muscle and liver of star-spotted dogfish (*Mustelus manazo*) from the northern region of Japan: A comparison with spiny dogfish (*Squalus acanthias*). *Arch. Environ. Contam. Toxicol.*, 64, 467-474.
4. Sakakibara, N., Hamasaki, T., Baba, M., Demizu, Y., Kurihara,



- M., Irie, K., Iwai, M., Asada, E., Kato, Y., and Maruyama, T. (2013). Synthesis and evaluation of novel 3-(3,5-dimethylbenzyl)uracil analogs as potential anti-HIV-1 agents. *Bioorg. Med. Chem.*, 21, 5900-5906.
5. Sakakibara, N., Tsukamoto, I., Isono, Y., Takata, M., Konishi, R., Kato, Y., and Maruyama, T. (2013). A new method for synthesis and angiogenic evaluation of leteprinim potassium and its novel analogs. *Heterocycles*, 87, 2369-2384.
 6. Umezawa, T., Ragamustari, S.K., Nakatsubo, T., Wada, S., Li, L., Yamamura, M., Sakakibara, N., Hattori, T., Suzuki, S., and Chiang, V.L. (2013). A lignan *O*-methyltransferase catalyzing the regioselective methylation of matairesinol in *Carthamus tinctorius*. *Plant Biotechnology*, 30, 97-109.
 7. Okabe, N., Nakamura, E., Himi, N., Narita, K., Tsukamoto, I., Maruyama, T., Sakakibara, N., Nakamura, T., Itano, T., and Miyamoto, O. (2013). Delayed administration of the nucleic acid analog 2Cl-C.OXT-A attenuates brain damage and enhances functional recovery after ischemic stroke. *Brain Research*, 1506, 115-131.
 8. Ohta, C., Haraguchi, K., Kato, Y., Endo, T., and Koga, N. (2013). Species difference in the metabolism of 2,2',3,4',5,5'-hexachlorobiphenyl (CB146) by animal and human liver microsomes. *Fukuoka Acta Medica* 104: 161-169.

[Others]

1. Ohta, C., Ogata, H., Yamamoto K., Haraguchi, K., Kato, Y., Endo, T., and Koga, N. (2016). *In vitro* metabolism of 5,7,3',4'-Tetramethoxyflavone by rat liver microsomes. *Bulletin of Nakamura Gakuen Univ and Nakamura Gakuen Univ Junior Coll* 48: 155-161.
2. Sakakibara, N. (2014). Synthesis and evaluation of novel nucleic acid derivatives as bioactive substances. *Yakugaku Zasshi*, 134, 965-972.
3. Sakakibara, N., Maruyama, T., and Kato, Y. (2013). Design and synthesis of novel nucleic acid analogs as potential angiogenesis or anti-HIV-1 agents. *Journal of Kagawa Pharmaceutical Association Kagayaku* 150, 68-70.
4. Ohta, C., Kato, Y., Haraguchi, K., Endo, T., and Koga, N. (2013). *In Vitro* metabolism of nobiletin in the small intestine and kidney of rats and guinea pigs. *Bulletin of Nakamura Gakuen Univ and Nakamura Gakuen Univ Junior Coll* 45, 141-149.

[Proceedings]

2016

1. Kato, Y., Haraguchi, K., Fujii, A., Fujii, Y., Kimura, O., Ohta, C., Endo, T., Koga, N., Yamada, S., Degawa, M. (2016). Induction of hepatic T₄ transporters by polychlorinated biphenyl in rats. *Organohalogen Compds* 78, 873-876.
2. Ohta, C., Fujii, Y., Haraguchi, K., Kato, Y., Kimura, O., Endo, T., and Koga, N. (2016). Metabolism of 2,2',3,4,4',5,6'-heptachlorobiphenyl (CB182) by rat, guinea pig and human liver microsomes. *Organohalogen Compds* 78, 870-872.

2014

1. Ohta, C., Haraguchi, K., Kato, Y., Endo, T., and Koga, N. (2014). The participation of rat CYP3A enzymes in the metabolism of 2,2',4,5,5'-pentachlorobiphenyl (CB101). *Organohalogen Compds* 76, 466-469.

2013

1. Ohta, C., Haraguchi, K., Kato, Y., Endo, T., Kimura, O., and Koga, N. (2013). *In vitro* metabolism of 2,2',4,4',5-pentachlorobiphenyl (CB99) by rat and guinea pig liver microsomes. *Organohalogen Compds* 75, 587-590.



Design, Manufacturing and Evaluation of Novel Pharmaceutical Preparations for Patients

Staff

Tadakazu Tokumura, Ph. D.

Professor since 2013

Previous position: Associate Professor of International University of Health and Welfare.

M.Sc. Graduated school of Agriculture, Kagawa University, 1981

Research

We have the research philosophy for Laboratory of Pharmaceutics, Kagawa School of Pharmaceutical Sciences, Tokushima Bunri University. Research projects were chosen based on the philosophy. The selected projects and those results were as follows:

- (1) Development of a novel dosage form containing fluticasone propionate for inflammatory bowel diseases

Crohn's disease and ulcerative colitis are the two primary types of inflammatory bowel diseases (IBD). Glucocorticosteroids were the standard treatment for IBD, but due to adverse events, their use was limited. However, new formulations of glucocorticosteroids have been developed to reduce systemic action. Fluticasone propionate (FLT) is an inhaled corticosteroid with high anti-inflammatory potency, used for the topical treatment of asthma. The purpose of this project was the design and preparation of a new FLT dosage form for topical treatment of IBD. The physicochemical property of FLT was very important for the dosage form design. The information for FLT was not enough, so we started the determination of the property. Chemical structures for degradation products of FLT in an alkaline solution were found by the support of Laboratory of Pharmacognosy and Natural Products Chemistry, which results were published. Basic studies for developing the new formulation were maintained.

- (2) Effect of the simple suspension method on the dissolution behavior

The simple suspension method is usually used. However, there was no report regarding the change of the dissolution property of the drug. The purpose of this project was a comparison of the dissolution profile between original and generic drugs after applied the simple suspension method. We evaluated the pharmaceutical preparations with valsartan and generic drugs of irbesartan.

- (3) Cleaning validation for machines used in the dispensary of pharmacy

When machines used in the dispensary, for an example, a

dividing and packing machine was applied for a granule or a powder, it was easily considered that the little amount of the granule or powder was left in the machine. Therefore, cleaning the machine was required. This cleaning will be performed according the procedure which is decided by each pharmacy. In the case of a pharmaceutical plant, cleaning validation was required for machines for manufacturing pharmaceutical preparations by GMP. The purpose of this project is to introduce the concept of cleaning validation to pharmacies. The preparations indicating the higher residual percent were researched on the basis of the result of questionnaire for pharmacists, which were the preparations of pranlukast hydrate, ketotifen fumarate, acetaminophen, and nicotinamide. The residual amounts of drugs on the machines were determined. In addition, cleaning effects of a cleaner with a dividing and packing machine, and cleaning using lactose on the amount of drugs in the machine were examined.

- (4) Degradation rate of olmesartan medoxomil in anaqueous solution

Degradation of olmesartan medoxomil in an aqueous solution and methanol was observed in the process for its determination method by HPLC. The degradation rates in aqueous solutions are determining. These results will provide the useful information for the application of the simple suspension method.

- (5) Development of novel pharmaceutical technology for poorly water-soluble drugs

To enhance oral bioavailability of poorly water-soluble drugs such as curcumin etc, we tried to prepare powders or water-based suspensions contain novel nano-particles by build-up or break-down methods, and estimated its particle characteristics. Increase of solubility of curcumin by γ -cyclodextrin was confirmed. The oral bioavailability was studied by rats.

- (6) Evaluation of powder preparations with sildenafil citrate for newborns in NICU

Sildenafil is a phosphodiesterase type 5 inhibitor that selectively reduces pulmonary vascular resistance in animal models and adult humans. Recent studies reported that the administration of sildenafil significantly increased oxygenation and reduced mortality with no clinically important side effects in infants with persistent pulmonary hypertension of the newborn.

A pharmaceutical preparation containing sildenafil citrate (SIL) for pulmonary arterial hypertension, Revatio Tablets 20 mg from Pfizer Japan Inc., is available for adults in Japan, whereas that for

children is not. Therefore, when sildenafil is administered to infants with persistent pulmonary hypertension of the newborn, a Revatio Tablets 20 mg is ground in a mortar to make a powder. Lactose is added to the powder as a diluent, and is mixed well in the mortar. The mixed powder is packaged for each dose using an automatic packaging machine. The contents of the drug in the packages were determined by HPLC. A part of this data was submitted and published. From this study the decrease of the content of sildenafil in the powder. A method for preventing the decrease was developed and reported.

(7) Development of new preparations with sildenafil for newborns in NICU

In the study for (6) the decrease of the content of sildenafil in the powder was found. A method for grinding tablets was developed, which was published. Furthermore, a method for preparing an oral liquid dosage form from Revatio Tablets was developed. These results were presented in an academic meeting.

(8) Determination of small particles to reach lung around a dividing and packing machine

It is easy to consider that there are small particles from dug preparations around dividing and packing machines. If small particles to reach lung are existed, safeties around people decrease. We examined using a cascade impactor. Small particles to reach lung were confirmed in pharmaceutical preparations of acetaminophen and theophylline.

Publications (2012~2017)

[Original papers]

2017

1. Tokumura T., Yoshida N., Mori-Yasumoto K., Shiota O., Kurita T., (2017). Degradation Rates and products of fluticasone propionate in alkaline solutions. *J Pharmaceutical Analysis* 7, 297-302.
2. Kawakami M., Kitada R., Kurita T., Tokumura T., (2017). A Method for Decreasing the Amount of the Drug Remaining on the Surfaces of the Mortar and Pestle after Grinding Small Amount of Tablets. *YAKUGAKU ZASSHI* 137(8), 1017-1025.

2016

3. Matsuyama S., Kurita T., and Tokumura T., (2016). Degradation Rate of Ebastine in an Aqueous Solution at pH 1.2 and the Effects of Cyclo dextrins. *Sch. Acad. J. Pharm.* 5(4), 87-91.
4. Tokumura T., Kawakami M., Kitada R. and Kurita T., (2016). Validated Assay Method for Fexofenadine Hydrochloride in Powder Preparations of Allegra® 60 mg Tablets to Develop a New Method for Grinding Tablets on Dispensing in Japan. *Sch. Acad. J. Pharm.* 5(8), 359-362.

2015

1. Kubodera M. Tokumura T. and Machida Y. (2015). Changes in the Amounts of Amoxicillin and Metronidazole Used for *Helicobacter pylori* Eradication Therapy in the Stomach after Their Oral Administration to Rats. *Sch. Acad. J. Pharm.* 4(3), 168-171.
2. Kino K., Nakatsuma A., Nochi H., Kiriya Y., Kurita T., Kobayashi T and Miyazawa H. (2015). Commentary on the Phototoxicity and Absorption of Vitamin B2 and Its

Degradation Product, Lumichrome (Commentary). *Pharm. Anal. Acta.*, 6: 403, doi: 10.4172/21532435.1000403

3. Tokumura T., Kawakami M., Kitada R., Yamamoto Hideki, Yamamoto Hiroshi, and Kurita T. (2015). Determination of Sildenafil Citrate in Powder Preparations Prepared from Revatio Tablets 20 mg for Infants with Persistent Pulmonary Hypertension of the Newborn. *Sch. Acad. J. Pharm.*, 4(8), 370-375.

2014

1. Tokumura T., Isaka, H., Kanou, M., Miyazaki, E., Kaneko, N., Kurita T., (2015). An inclusion complex of fluticasone propionate with γ -cyclodextrin in aqueous solution and in a solid state. *J. Drug Del. Sci. Tech.* 26, 24-27.
2. Tokumura T., Kanou, M., Miyazaki, E., Kaneko, N., Isaka, H. (2014). Degradation rate of fluticasone propionate in an alkaline solution of 0.1N NaOH : methanol = 1 : 1. *Int. Res. J. Pharm. App. Sci.*, 4(5), 1-3.
3. Tokumura T., Miyazaki, E., Isaka, H., Kaneko, N., Kanou, M., (2014). Solubility of fluticasone propionate in aqueous solutions measured by a method avoiding its adsorption to experimental tools. *Int. Res. J. App. Sci.*, 4(4), 19-24.

2013

1. Kurita T., Makino, Y. (2013). Novel curcumin oral delivery systems. *Anticancer Res* 33, 2807-2821.

2012

1. Kubodera, M., Tokumura T., and Machida, Y., (2012). Determination of metronidazole in a rat stomach by HPLC for obtaining basic data of eradication therapy of *Helicobacter pylori*. *J Pharmaceutical Analysis* 2, 378-381.
2. Tokumura T., Nagaoka, M., and Machida, Y., (2012). Effect of doses and dosage forms on the bioavailability of amoxicillin in non-fasted rats. *J Drug Del Sci Tech* 22, 568-570.



Laboratory of Pharmaceutical Health Care and Sciences

Staff

Masaki Ninomiva, Ph. D.

Professor since 2008

Doctor of medical science, University of Kagawa, 1995

Naomi Iihara, Ph.D.

Professor since 2011

Ph.D. University of Okayama; Pharmacist

Hiroaki Ikeda, Ph.D.

Professor since 2016

Ph.D. University of Hiroshima; Pharmacist

Akira Nakatsuma, Ph. D.

Lecturer since 2017

Ph. D. University of Okayama, 2001

Taketo Okada, Ph.D.

Assistant Professor since 2013

Ph.D. Graduate School of Medical and Pharmaceutical Sciences,
Chiba University; Pharmacist

Research

(1) Patient–Healthcare Professional Relationship

Patients accept or refuse their medication therapy based on their personal beliefs. We analyze relationship between patient's perceptions of medication therapy and their behavior such as medication adherence.

We developed Medication Acceptance, Preference and Adherence Scale (MAPAS), which assessed each patient's beliefs, values and ideas concerning their acceptance and preference for medications and treatments. We found that patients' dissatisfaction consistent determinants of intentional non-adherence to medication, but not unintentional non-adherence. In addition, we found that cancer patients prefer aggressive therapies, even when self-estimations of ADR endurance are not very high, especially if they have been receiving chemotherapy for a short period of time.

(2) Pharmacoepidemiology Study Encouraging Proper Use of Medication

Post-marketing surveillance is important because it can evaluate medication use in the real world.

We surveyed use of medication with driving with prohibitions or cautions in outpatient settings for patients aged 25 years and older using the National Health Insurance Claims Database. Of outpatients aged 25 years and older who were administered medications, 73% outpatients were given the medications with

cautions or prohibitions on driving. For the elderly, prescriptions were found with dosages that often exceeded the recommended limits.

(3) Chem- and Bio-Informatics of Traditional Medication Systems, and Their Related Medicinal Resources

We perform the informatics and computational science based on evidences related to medical and pharmaceutical issues. This research has been mainly demonstrated by chemical and biological experiments, database construction, and multivariate statistical analysis. In recent studies, we have focused on the theoretical analysis of a traditional and an empirical medication system in *Kampo* (traditional Japanese medicine), and the transcriptome and metabolome analyses of medicinal resources related to *Kampo*.

(4) Modulation of multi-drug resistance related protein transport by interaction with dietary supplements

An interaction is taken to be the situation in which administration of a drug or substance induces changes in the pharmacokinetics of another simultaneously administered drug – by increasing either the plasma or intracellular concentration of the latter, and thus giving rise to the possibility of an adverse reaction.

The ABC-transporter superfamily, which functions as a drug efflux pump, is known to limit the absorption of a variety of drugs. We investigated the effects of food extracts on anticancer drug transport by the multi-drug resistance related proteins (MRPs). MRPs are efflux transporters expressed in human glioblastoma cell line T98G. The effects on MRP mediated transport were also evaluated using calcein, which is the substrate of MRP. Acute exposure to kaempferol caused a concentration-dependent decrease in the extracellular efflux of calcein compared with the control. As for the simultaneous exposure to kaempferol and cisplatin, the cytotoxicity of cisplatin was expected to be potent because MRP and glutathione *S*-transferases (GST) are both inhibited by kaempferol. However, the cytotoxicity of cisplatin decreased.

Western blot analysis and reverse transcription–polymerase chain reaction (RT–PCR) showed that treatment with 10 and 20 μ M kaempferol for up to 72 hr was able to significantly lower MRP2 expression, whereas increased expression in a concentration-dependent on GST mRNA and protein levels. Furthermore, GST was strongly activated in T98G cell treated with kaempferol.

The results of the study also point to possible kaempferol-drug interaction, especially when the cytotoxicity of anticancer drugs are dependent on glutathione *S*-transferases and MRP-mediated transport processes. Hereafter, these possible efficacies need to be examined under in vivo conditions in detail.

Publications

* 2013-2017

[Original papers]

2017

1. Iihara, N., Ohara, E., Nishio, T., Muguruma, H., Matsuoka, E., Houchi, H., and Kirino, Y. (2017) Patient Preference for Aggressive Medication Therapies with Potentially Stronger Adverse Drug Reactions Revealed Using a Scenario-based Survey YAKUGAKU ZASSHI 137(9), 1161-1167

2016

1. Iihara N., Bando Y, Ohara M, Yoshida T, Nishio T, Okada T., and Kirino Y (2016) Polypharmacy of medications and fall-related fractures in older people in Japan: a comparison between driving-prohibited and driving-cautioned medications. J Clin Pharm Ther 41(3), 273—278
2. Okada T., Afendi FM, Yamazaki M, Chida KN, Suzuki M, Kawai R, Kim M, Namiki T, Kanaya S, and Saito K (2016) Informatics framework of traditional Sino-Japanese medicine (Kampo) unveiled by factor analysis. J Nat Med 70(1), 107—114
3. Watabe, S., Morikawa, M., Kaneda, M., Nakaishi, K., Nakatsuma, A., Ninomiya, M., Yoshimura, T., Miura, T., Ito, E. (2016) Ultrasensitive detection of proteins and sugars at single-cell level. Commun Integr Biol. 9(1) e1124201
4. Okada T., Takahashi H, Suzuki Y, Sugano S, Noji M, Kenmoku H, Toyota M, Kanaya S, Kawahara N, Asakawa Y, and Sekita S (2016) Comparative analysis of transcriptomes in aerial stems and roots of *Ephedra sinica* based on high-throughput mRNA sequencing. Genom Data 10, 4—11

2015

1. Iihara, N., Nishio, T., Goda, M., Anzai, H., Kagawa, M., Houchi, H., and Kirino, Y. (2015). Effect of endurance for adverse drug reactions on the preference for aggressive treatments in cancer patients. Support Care Cancer 23(4), 1091-1097.
2. Nakatsuma, A., Kiriya, Y., Kino, K., Ninomiya, M. (2015) Diabetes drugs that protect pancreatic β cells. Integr. Mol. Med. 3(1) .
3. Nakatsuma, A., Kaneda, M., Kodama, H., Morikawa, M., Watabe, S., Nakaishi, K., Yamashita, M., Yoshimura, T., Miura, T., Ninomiya, M., Ito, E. (2015) Detection of HIV-1 p24 at Attomole Level by Ultrasensitive ELISA with Thio-NAD Cycling. PLoS One. 10(6):e0131319.
4. Kino, K., Nakatsuma, A., Nochi, H., Kiriya, Y., Kurita, T., Kobayashi, T., Miyazawa, H. (2015) Commentary on the Phototoxicity and Absorption of Vitamin B2 and Its Degradation Product, Lumichrome. Pharm Anal Acta 6:403.
5. Nakatsuma, A., Kaneda, M., Kodama, H., Morikawa, M., Watabe, A., Nakaishi, K., Yamashita, M., Yoshimura, T., Miura, T., Ninomiya, M., Ito, E. (2015) Ultrasensitive colorimetric detection of HIV-1 p24. Clinical Laboratory International, 2015 October Issue, 20-25.
6. Nakatsuma, A., Wada, S., Kamano, J., Kiriya, Y., Kino, K., Ninomiya, M. (2015) The effects of herbal teas on drug permeability. Integr. Mol. Med. 3(1) : 453-456.

2014

1. Iihara N., Yoshida T, Okada T., Nakatsuma A., and Kirino Y (2014) Survey of usage of medication with driving with prohibition or caution by the national health insurance claims database in Japan. Jpn J Pharm Health Care Sci 40(2), 67—77

2013

1. Iihara, N., Nishio, T., Okura, M., Anzai, H., Kagawa, M., Houchi, H., and Kirino, Y. (2013). Comparing patient dissatisfaction and rational judgment in intentional medication non-adherence versus unintentional non-adherence. J Clin Pharm Ther 39 (1), 45-52.

[Review articles]

1. Iihara, N. (2014). “A Continuing Education Program for Pharmacists to Assess Adverse Drug Reactions” Journal of Pharmaceutical Science and Technology 74(5), 298-300.
2. Tsuchiya, F., and Iihara, N. (2014). “Healthcare IT and medication: How does healthcare IT-ization affect medication development and medication safety assurance?” YAKUGAKU ZASSHI 134(5), 583-584.
3. Iihara, N., and Kirino, Y. (2014). “A community electronic prescription system connecting physicians, pharmacists, and patients, and utilization of clinical information” YAKUGAKU ZASSHI 134(5), 589-593.

[Books]

1. Okada T., and Noji M (2014) Metabolome analysis of medicinal plants and crude drugs –a study case of *Ephedra* plants by comprehensive metabolite analysis using mass spectrometry and multivariate statistical analysis– (Chapter 1 in Part III). In: Kawahara N (Sv) Recent progress of studies on medicinal plants and crude drugs –cultivation and quality evaluation of medicinal plants, and development of Kampo products–. CMC Publishing, Tokyo, pp 122—131 (in Japanese)
2. Okada T., Afendi FM, Katoh A, Hirai A, and Kanaya S (2013) Multivariate analysis of analytical chemistry data and utility of the KNAPSAcK family database to understand metabolic diversity in medicinal plants (Chapter 18). In: Chandra S, Lata H, and Varma A (Eds) Biotechnology for medicinal plants: micropropagation and improvement. Springer, Berlin Heidelberg, pp 413—438



Laboratory of Pharmaceutical Education

Staff

Hiroshi Tokumaru, Ph. D.

Professor since 2012

Research Assistant Professor, Dpt. Neurobiology, Duke University,
NC, U.S.A.

D.Sc. Kyushu University, Pharmaceutical Sciences, 1989

Takayuki Ohshima, Ph. D.

Associate Professor since 2008

Ph.D. Graduate School of Agriculture, Tsukuba University, 2001

Takaaki Shirahata, Ph.D.

Lecturer since 2013

Ph.D. Graduate school of Pharmaceutical Sciences, University of
Tokyo, 2006

Hisayo Sadamoto, Ph. D.

Lecturer since 2014

Assistant Professor since 2005

Ph. D. in Hokkaido University, Biological Sciences, 2002

Suguru Kobayashi, Ph. D.

Assistant Professor since 2005

Assistant Professor, Sapporo Medical University

Ph. D. in Hokkaido University, Biological Sciences, 2000

Risa Mukai, Ph. D.

Postdoctoral Scholar since 2015

Ph.D. Graduate School of Engineering, Tokushima Bunri
University, 2015

Research

Theme 1. The molecular mechanism of complexin (Hiroshi Tokumaru, Hisayo Sadamoto)

Action potential-evoked neurotransmitter release is triggered by Ca^{2+} influx through voltage-gated calcium channels located next to the active zone. The increase in Ca^{2+} concentration initiates rapid signaling cascades that lead to the exocytosis of synaptic vesicles containing high concentrations of neurotransmitter. Two soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) proteins from the presynaptic membrane, syntaxin 1 and SNAP-25, and one SNARE protein from the synaptic vesicle membrane, synaptobrevin-2 (also known as VAMP-2), form a

four-helix bundle (called the *trans*-SNARE complex or SNAREpin) that catalyzes membrane fusion. The synaptic vesicle protein synaptotagmin 1 (Syt1) serves as a major Ca^{2+} sensor for fast action potential-evoked synaptic vesicle exocytosis. The rapid interactions between Syt1, the SNARE complex and membrane phospholipids induced by Ca^{2+} are critical for membrane fusion.

The precise control of evoked neurotransmitter release requires several cytosolic proteins including complexin (also known as synaphin). Complexin and its binding to the SNARE complex are critical for fast neurotransmitter release, as demonstrated by studies in knockout mice and in *Drosophila* mutants. In addition, intra-presynaptic terminal injection of a SNARE-binding domain peptide that blocks complexin binding to the SNARE complex also inhibits rapid neurotransmitter release. Despite this evidence, complexin's function remains controversial. Furthermore, recent studies suggest that complexin contains several functional domains that either stimulate or inhibit neurotransmitter release. Thus, the function of complexin is likely more complex than expected from its small size (134 aa for mammalian complexin 1 and 2).

Like Syt1-deficient mice, complexin 1/2 double-knockouts exhibit an impairment of action potential-evoked synchronous (but not asynchronous) neurotransmitter release. However, an important difference exists between the two types of knockout mice: elevated external Ca^{2+} can rescue synchronous release in complexin 1/2 double-knockouts but not in Syt1 knockouts. Thus, the functions of complexin and Syt1 in action potential-evoked fast neurotransmitter release are intimately related to each other yet distinct.

Biochemically investigating the relationship between the two proteins, we previously demonstrated that complexin directly binds to Syt1 even in the absence of Ca^{2+} . Because Syt1 binds to SNARE complexes weakly in the absence of Ca^{2+} , we proposed that complexin recruits Syt1 to the SNARE complex prior to Ca^{2+} influx. Recently, we examined the interaction between complexin, Syt1 and the SNARE complex using recombinant proteins. Our results indicate that Syt1 recruitment to the SNARE-driven fusion machinery by complexin is essential for vesicle exocytosis.

Syt1 bound weakly to complexin alone, but the addition of three SNARE proteins (syntaxin 1, SNAP-25 and VAMP/synaptobrevin-2) in combination, but not individually, markedly enhanced binding. Unlike full-length complexin (amino acid [aa] 1–134) and an NH₂ (N)-terminally truncated complexin (aa 46–134), carboxy (C)-terminally truncated complexin s (aa 1–104 and 1–124) could not support Syt1 binding even in the presence of the SNARE complex. These results indicate that the binding of Syt1 to the C-terminal region of complexin promotes its

Laboratory of Pharmaceutical Education

recruitment to the SNARE-driven fusion machinery, and that this process is crucial for Ca^{2+} -dependent vesicle exocytosis.

Theme 2. Differential localizations of GKAP/SAPAP1 isoforms in developing hippocampal neurons (Hisayo Sadamoto)

Guanylate kinase-associated protein (GKAP) and SAP90/PSD-95-associated protein 1 (SAPAP1) proteins form complexes with PSD-95 and Shank at excitatory postsynaptic sites, and are implicated in synapse formation and synaptic plasticity. GKAP/SAPAP1 proteins, which displayed different N termini, have appeared as multiple alternatively spliced isoforms. However, specific functional roles of individual isoforms still remain unclear. To understand particular functions of GKAP/SAPAP1 isoforms in formation and maintenance of synaptic connections, we here investigated expression and subcellular distributions of these isoforms in hippocampal neurons during synaptic development. First, we identified two isoforms of SAPAP1 (named as SAPAP1b and 1c) in mice hippocampus, which exhibited an alternative usage of two exons in the middle part of SAPAP1 transcript. Using primary culture of mouse hippocampal neurons and confocal microscope, we sought to examine localizations of each EGFP-tagged GKAP/SAPAP1 isoform (GKAP, SAPAP1, SAPAP1b or SAPAP1c). During synaptic maturation, GKAP/SAPAP1 isoforms were found to display differences in cluster formation at the dendritic spines. EGFP-SAPAP1 formed clusters at dendritic shafts on an early stage of synapse formation and did not change the rate of accumulation (clustering index) in mature dendritic spines at later stages. Clusters of EGFP-SAPAP1b and EGFP-SAPAP1c were also found to occur at an early stage, but tended to disappear during synaptic maturation. In contrast, GKAP clearly accumulated in dendritic spines at a later stage of synaptic maturation. These results suggest the possibility that each spliced isoform of GKAP/SAPAP1 has a specific function in synapse formation.

Theme 3. Synaptic modulation and oscillatory network modulation in odor information processing (Suguru Kobayashi)

Synchronous oscillatory activity is common in the olfactory behavior of both vertebrates and invertebrates. In the olfactory center of terrestrial animals, changes in the oscillatory frequency of the local field potential (LFP) are thought to be involved in olfaction-based behavior and olfactory memory. We study GABAergic and FMRFamideergic neuromodulation of oscillatory activity in odor information processing of the procerebrum (PC) in the land slug *Limax valentianus*. We found that GABA and FMRFamide are present in the PC and these modulatory roles are involved in the oscillatory neural network of the PC. A part of results for excitatory GABAergic and FMRFamideergic

neuromodulation are published in *J. Neurophysiol.* (2012) and *Eur. J. Neurosci.* (2010). Recent study showed the presence of cholinergic excitatory modulation for PC neurons via nicotinic ACh receptors activation (*J Comp Neurol*, 2014) and feedforward inhibition in the cholinergic afferents from the tentacles to the PC. Furthermore, in comprehensive analysis with *in vitro* cultured PC networks, the presence and role of biogenic amines were discussed (*J Comp Neurol*, 2016). We use electrophysiology and optical recording methods to understand the role of oscillatory dynamics in odor recognition and memory storage.

Grant Support: Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research.

Collaborations: With other universities.

Theme 4. Functional regulation of proteins by post-translational modifications (Ohshima)

Post-translational modifications such as ubiquitination, phosphorylation, and acetylation regulate the function of many proteins. Recently, a number of ubiquitin-like proteins (Ubl) have been identified that are covalently linked to lysine residues in target proteins. One Ubl, SUMO-1, also known as PIC1, UBL1, sentrin, GMP1, and SMT3, is an 11-kDa protein that is structurally homologous to ubiquitin. SUMO-1 modification plays an important role in altering the function of modified proteins, including transcriptional activation, nuclear localization, and decreased turnover. SUMO-1 is conjugated to proteins through a series of enzymatic steps. Initially, the ATP-dependent formation of a thioester bond between SUMO-1 and the E1 enzyme complex (SAE1/Uba2) is formed, and SUMO-1 is then transferred to the E2-conjugating enzyme Ubc9. Finally, SUMO-1 is conjugated from Ubc9 directly to a lysine residue of target proteins. The E3 ligase that conjugates SUMO-1 to target molecules *in vitro* and *in vivo* has only recently been identified. One group of such E3 ligases, protein inhibitor of activated STAT (PIAS) family proteins homologous to the yeast Siz family protein, has a conserved RING-finger domain, which regulates transactivation of many transcription factors by conjugating SUMO-1. In order to understand the molecular mechanisms by which transcriptional regulation through SUMO-1 modification, we focus the transcription (co)factors involving in cell growth, differentiation, immortalization and attempt to define the biological significance of sumoylation in carcinogenesis.

Theme 5. Analyzing the molecular mechanisms by which human T-cell leukemia virus type-1 (HTLV-1) infection is the cause of morbidity and mortality in adult T-cell leukemia (ATL) (Mukai and Ohshima).

Human T-cell leukemia virus type 1 (HTLV-1), a human retrovirus, is the causative agent of adult T-cell leukemia (ATL), an aggressive malignancy of CD4^+ T lymphocytes. HTLV-1 is also closely related to HTLV-1-associated myelopathy or tropical



spastic paraparesis (HAM/TSP). Although most HTLV-1-infected people are non-symptomatic, approximately 4% of patients develop ATL after a long period of latent infection (over 50 years). To date, there are few effective therapies available for ATL patients, possibly due to a lack of detailed information about the molecular mechanism of cell growth regulation by HTLV-1.

HTLV-1 basic leucine-zipper factor, HBZ, which is encoded in the complementary strand of the HTLV-1 genome, has been identified. HBZ is a nuclear protein that contains a transactivation domain and a basic leucine-zipper (bZIP) domain in its N- and C-termini, respectively. HBZ interacts with cellular bZIP proteins, in particular with the AP-1 family of transcription factors, and regulates their transcriptional activities, resulting in the control of viral gene transcription from the HTLV-1 promoter. In contrast to other viral protein, HBZ is constitutively expressed in all ATL patient samples because its 3'-LTR is conserved and unmethylated in ATL cells. HBZ may play a central role in the pathogenesis of ATL, however, its function has not yet been understood.

Theme 6. Analysis of Nonlinear Phenomena of Mathematical Models of Neurons (Takaaki Shirahata)

Individual neurons are capable of producing action potentials or spikes in response to external stimulation through the interactions of various ionic channels expressed on the plasma membrane of neurons. Based on data derived from the electrophysiological experiments, various mathematical models which reproduce the electrical excitability of individual neurons are developed (e.g., nonlinear ordinary differential equations). By analyzing the mathematical models in detail one can understand the dynamics of individual neurons. Modeling studies are suggested to be useful for studying drugs which are used for the therapy of channelopathy. In addition, based on the theory proposed by US mathematician and physicist, modeling studies are also expected to be useful for developing medical devices. Taking the above into account, this laboratory is investigating ordinary differential equations which describe the electrical excitability of individual neurons.

1. Research about a ghostbursting phenomenon of electrosensory pyramidal neurons in weakly electric fish
2. Research about pacemaking of RPa1 neurons in snail *Helix pomatia*
3. Research about the spiking activity of neocortical pyramidal neurons
4. Research about the spiking activity of vibrissa motoneurons
5. Research about the spiking activity of circadian pacemaker neurons in the suprachiasmatic nucleus
6. Research about the spiking activity of medial vestibular nucleus neurons
7. Research about the dynamics of retinal horizontal cells

8. Research about the dynamics of pseudo-plateau bursting of endocrine cells

Publications (2012~2017)

[Original papers]

2017

1. Shirahata, T. (2017). Dependence of the ghostbursting model's dynamical states on the current injected into the dendritic compartment and the ratio of somatic to total surface areas. *Advanced Studies in Theoretical Physics* 11, 561-565.
2. Shirahata, T. (2017). A Mathematical Modeling Study of Dopaminergic Retinal Neurons under Hyperpolarized Conditions. *International Journal of Theoretical and Mathematical Physics* 7, 4-8.

2016

3. Matsuo R., Tanaka M., Fukata R., Kobayashi S., Aonuma H., Matsuo Y. (2016) Octopaminergic system in the central nervous system of the terrestrial slug *Limax*. *J Comp Neurol* 524, 3849-3864.
4. Matsuo R., Fukata R., Kumagai M., Kobayashi A., Kobayashi S., Matsuo Y. (2016) Distribution of histaminergic neurons and their modulatory effects on oscillatory activity in the olfactory center of the terrestrial slug *Limax*. *J Comp Neurol* 524, 119-35.
5. Mukai R. and Ohshima T. (2016) Enhanced stabilization of MCL1 by the human T-cell leukemia virus 1 bZIP factor is modulated by blocking the recruitment of cullin 1 to the SCF complex. *Mol. Cell. Biol.* 36, 3075-3085.
6. Klionsky, D. J., et al (including Ohshima T.). (2016) Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). *Autophagy* 12, 1-222.
7. Shirahata, T. (2016). The Relationship of Sodium and Potassium Conductances with Dynamic States of a Mathematical Model of Electrosensory Pyramidal Neurons. *Applied Mathematics* 7(9), 819-823.
8. Shirahata, T. (2016). Dynamics of a Pituitary Cell Model: Dependence on Long-Lasting External Stimulation and Potassium Conductance Kinetics. *Applied Mathematics* 7(9), 861-866.
9. Shirahata, T. (2016). Quantitative evaluations of the contribution of the excitatory ionic conductance to repetitive spiking in a mathematical model of medial vestibular nucleus neurons. *Acta Biologica Hungarica* 67(2), 215-219.
10. Shirahata, T. (2016). Dynamic Behavior Induced by the Cooperation between Two Different Ionic Conductances in a Mathematical Model of Vibrissa Motoneurons. *Applied Mathematics* 7(10), 1043-1048.
11. Shirahata, T. (2016). The Effect of Variations in Ionic Conductance Values on the Suppression of Repetitive Spiking in a Mathematical Model of Type-A Medial Vestibular Nucleus Neurons. *Applied Mathematics* 7(10), 1134-1139.
12. Shirahata, T. (2016). Evaluating Bistability in a Mathematical Model of Circadian Pacemaker Neurons. *International Journal of Theoretical and Mathematical Physics* 6(3), 99-103.
13. Shirahata, T. (2016). The Effect of Variations in Ionic Conductance Values on the Dynamics of a Mathematical Model of Non-Spiking A-Type Horizontal Cells in the Rabbit Retina. *Applied Mathematics* 7(12), 1297-1302.

2015

14. Takahashi N., Sawada W., Noguchi J., Watanabe S., Ucar H.,

Laboratory of Pharmaceutical Education

- Hayashi-Takagi A., Yagishita S., Ohno M., Tokumaru H. and Kasai H. Two-photon fluorescence lifetime imaging of primed SNARE complexes in presynaptic terminals and β cells. *Nature Commun* 6, 1-15.
15. Mukai R., and Ohshima T. (2016) HTLV-1 bZIP factor suppresses the centromere protein B (CENP-B)-mediated trimethylation of histone H3K9 through the abrogation of DNA-binding ability of CENP-B. *J. Gen. Virol.* 96, 159-164.
16. Watanabe S., Takanashi F., Ishida K., Kobayashi S., Kitamura Y., Hamasaki Y., Saito M. (2015) Nitric Oxide-Mediated Modulation of Central Network Dynamics during Olfactory Perception. *PLoS One* 10, e0136846.
17. Shirahata T. (2015). Evaluation of kinetic properties of dendritic potassium current in ghostbursting model of electrosensory neurons. *Applied Mathematics* 6(1), 128-135.
18. Shirahata T. (2015). Numerical Simulation Analysis of a Mathematical Model of Circadian Pacemaker Neurons. *Applied Mathematics* 6(8), 1214-1219.
19. Shirahata T. (2015). Numerical Study of a Mathematical Model of Vibrissa Motoneurons: The Relationship between Repetitive Spiking and Two Types of Sodium Conductance. *International Journal of Theoretical and Mathematical Physics* 5(3), 48-52.
20. Shirahata T. (2015). Numerical Simulation of Bistability between Regular Bursting and Chaotic Spiking in a Mathematical Model of Snail Neurons. *International Journal of Theoretical and Mathematical Physics* 5(5), 145-150.
- 2014**
21. Mukai R., and Ohshima T. (2014) HTLV-1 HBZ positively regulates the mTOR signaling pathway via inhibition of GADD34 activity in the cytoplasm. *Oncogene* 33, 2317-2328.
22. Toyama M., Aoyama H., Mukai R., Nakamura M., Yoshimura K., Okamoto M., Ohshima T., Hashimoto Y., and Baba M. (2014) A novel tetramethylnaphthalene derivative selectively inhibits adult T-cell leukemia (ATL) cells in vitro. *Anticancer Res.* 34, 1771-1778.
23. Matsuo R., Kobayashi S., Wakiya K., Yamagishi M., Fukuoka M., Ito E. (2014) The cholinergic system in the olfactory center of the terrestrial slug *Limax*. *J Comp Neurol* 522, 2951-2966.
24. Shirahata T. (2014). Effect of sodium conductance variations on electrical behavior of a neocortical neuron model. *Acta Biologica Hungarica* 65(4), 379-384.
- 2013**
25. Watanabe T., Sadamoto H., and Aonuma H. (2013) Molecular basis of the dopaminergic system in the cricket *Gryllus bimaculatus*. *Invert Neurosci* 13, 107-23.
26. Sadamoto H. and Muto H. (2013) Fluorescence Cross-correlation Spectroscopy (FCCS) to Observe Dimerization of Transcription Factors in Living Cells. *Methods Mol Biol* 977, 229-41.
27. Murakami J., Okada R., Sadamoto H., Kobayashi S., Mita K., Sakamoto Y., Yamagishi M., Hatakeyama D., Otsuka E., Okuta A., Sunada H., Takigami S., Sakakibara M., Fujito Y., Awaji M., Moriyama S., Lukowiak K. and Ito E. (2013) Involvement of insulin-like peptide in long-term synaptic plasticity and long-term memory of the pond snail *Lymnaea stagnalis*. *J Neurosci* 33, 371-83.
28. Shirahata T. (2013). Novel types of bistability in a model of a bursting pacemaker neuron RPa1 from the snail, *Helix pomatia*. *Acta Biologica Hungarica* 64(1), 131-135.
- 2012**
29. Torikoshi K., Abe H., Matsubara T., Hirano T., Ohshima T., Murakami T., Araki M., Mima A., Iehara N., Fukatsu A., Kita T., Arai H., and Doi T. (2012) Protein inhibitor of activated STAT, PIASy, regulates alpha-smooth muscle actin expression by interacting with E12 in mesangial cells. *PLoS One* 7, e41186-41199.
30. Kobayashi S., Ito E. (2012) GABAergic effects on the slow oscillatory neural activities in the procererebrum of *Limax valentianus*. *Acta Biologica Hungarica* 63 (Suppl. 2), pp. 217-221.
31. Elekes K., Battonyai I., Kobayashi S., Ito E. (2012) Organization of the procererebrum in terrestrial pulmonates (*Helix*, *Limax*) reconsidered: cell mass layer synaptology and its serotonergic input system. *Brain Structure and Function* 218, 477-490.
32. Kobayashi S., Matsuo, R., Sadamoto H., Watanabe, S., and Ito, E. (2012) Excitatory effects of GABA on procererebrum neurons in a slug. *J Neurophysiol* 108, 989-998.
33. Sadamoto H., Takahashi, H., Okada, T., Kenmoku, H., Toyota, M., Asakawa, Y. (2012) De novo sequencing and transcriptome analysis of the central nervous system of mollusc *Lymnaea stagnalis* by deep RNA sequencing. *PLoS One* 7 e42546.
34. Ito E., Otsuka E., Hama N., Aonuma H., Okada R., Hatakeyama D., Fujito Y., Kobayashi S. (2012) Memory trace in feeding neural circuitry underlying conditioned taste aversion in *Lymnaea*. *PLoS One* 7 e43151.
35. Shirahata T. (2012). Analysis of the electrosensory pyramidal cell bursting model for weakly electric fish: Model prediction under low levels of dendritic potassium conductance. *Acta Biologica Hungarica* 63(3), 313-320.

[Books]

- Mochida S., Tokumaru H. et al., (2015) "Presynaptic Terminals"
Springer ISBN978-4-431-55165-2



Center for Instrumental Analysis

Staff

Professor: Kentaro Yamaguchi, Ph. D (Apr. 2004) (LAC)

Associate Professor: Hajime Takeuchi, Ph.D. (Apr. 2013) (LPHS)

Lecturer: Kazuaki Ohara, D. Eng. (Apr. 2010)

Educational History:

Graduated from Graduate School of Tokyo University
in Mar. 1992

Research

Observation of the giant molecules by means of mass spectrometry:

Mass Spectrometry (MS) has been developed and adopted to wide variety of analytical chemistry in recent years.

Although MS was basically developed for high molecular weight substances in the field of biochemistry, the measurement of huge molecules over 10k Da is still very difficult. This is caused by the ionizing problems, stability of the compounds and the existence of various impurities.

We develop some new techniques to overcome these problems by using newly equipped FT-ICR mass spectrometer.

Crystalline Sponge-Laser Desorption Ionization:

Recently, crystalline sponge (CS) method was discovered to analyze non-crystalline compounds by means of X-ray crystallography. The laser desorption ionization (LDI) mass spectrometry (MS) is first adopted to use this CS as a matrix for ionization. Since then, this new ionization method (CS-LDI MS) has been developed in this laboratory.

Publications (2013~2018)

[Original papers]

2018

- Hayashi, Y.; [Ohara, K.](#); Taki, R.; Saeki, T.; *Yamaguchi, K. (2018). Combined analysis of 1,3-benzodioxoles by crystalline sponge X-ray crystallography and laser desorption ionization mass spectrometry. *Analyst*, 2018, 143, 1475-1481.
- Hirao, T.; Kim, D-S.; Chi, X.; Lynch, V-M., [Ohara, K.](#); *Park, J-S.; *Yamaguchi, K.; *Sessler, J-L. (2018). Control over multiple molecular states with directional changes driven by molecular recognition. *Nat. Commun.*, 2018, 9, 823.
- *Tominaga, M.; Ando, H.; [Ohara, K.](#); Itho, T.; Yamaguchi, K. (2018). Crystal Formation of a Coordination Cage through Spherical Particles Derived from a Tripodal Ligand Containing Adamantane Moiety and Silver(I) Salt. *Chem. Lett.*, 2018, 47, 315-317.

2017

- *Tominaga, M.; Takahashi, E.; Ukai, H.; [Ohara, K.](#); Itho, T.; Yamaguchi, K. (2017). Solvent-Dependent Self-Assembly and Crystal Structures of a Salen-Based Macrocyclic. *Org. Lett.*, 2017, 19(7), 1508-1511.

2016

- *Tominaga, M.; Kawaguchi, T.; [Ohara, K.](#); Yamaguchi, K.; Katagiri, K.; Itho, T.; *Azumaya, I. (2016). Vesicle Formation of Three-dimensional Trinuclear Silver(I) Complexes Built from Tris-NHC Ligands Bearing Long Alkyl Chains. *Chem. Lett.*, 2016, 45, 1201-1203.
- *Tominaga, M.; Noda, A.; [Ohara, K.](#); Yamaguchi, K.; Itho, T. (2016). Synthesis, Hollow Spherical Aggregation, and Crystallization of an Adamantane-derived Azacyclophane Containing Triazine Rings. *Chem. Lett.*, 2016, 45, 733-775.
- *[Ohara, K.](#); Tominaga, M.; Masu, H.; Azumaya, I.; *Yamaguchi, K. (2016). Adamantane-based Bidentate Metal Complexes in Crystalline and Solution State. *Anal. Sci.*, 2016, 32(12), 1347-1352.
- *Kawahata, M.; Komagawa, S.; [Ohara, K.](#); Fujita, M.; *Yamaguchi, K. (2016). High-resolution X-ray structure of methyl salicylate, a time-honored oily medicinal drug, solved by crystalline sponge method. *Tetrahedron Lett.*, 2016, 57, 4633-4636.
- Sawada, T.; Yamagami, M.; [Ohara, K.](#); Yamaguchi, K.; *Fujita, M. (2016). Peptide [4]Catenane by Folding and Assembly. *Angew. Chem. Int. Ed.*, 2016, 55, 4519-4522.
- Ishizuka, T.; Watanabe, A.; Kotani, H.; Hong, D.; Satonaka, K.; Wada, T.; Shiota, Y.; Yoshizawa, K.; [Ohara, K.](#); Yamaguchi, K.; Kato, S.; *Fukuzumi, S.; *Kojima, T. (2016). Homogeneous Photocatalytic Water Oxidation with a Dinuclear Co^{III}-Pyridylmethylamine Complex. *Inorg. Chem.*, 55, 1154-1164.
- Wang, S.; Sawada, T.; [Ohara, K.](#); Yamaguchi, K.; *Fujita, M. (2016). Capsule-Capsule Conversion by Guest Encapsulation. *Angew. Chem. Int. Ed.*, 55, 2063-2066.
- *Tominaga, M.; Kawaguchi, T.; [Ohara, K.](#); Yamaguchi, K.; Masu, H.; *Azumaya, I. (2016). Synthesis and crystal structures of twisted three-dimensional assemblies of adamantane-bridged tris-NHC ligands and Ag^I. *CrystEngComm*, 18, 266-273.

2015

- *Ikeda, A.; Iwata, N.; Hino, S.; Mae, T.; Tsuchiya, Y.;

Center for Instrumental Analysis

- Sugikawa, K.; Hirao, T.; Haino, T.; Ohara, K.;
Yamaguchi, K. (2015). Liposome collapse resulting from an
allosteric interaction between 2,6-dimethyl- β -cyclodextrins
and lipids. *RSC Adv.*, 5, 77746-77754.
2. Ohara, K.; Nakai, A.; Yamaguchi, K. (2015). Laser desorption
ionization of stilbenes in crystalline sponge.
Eur. J. Mass Spectrom., 21, 413-421.
4. Shinozaki, Y.; Richard G.; Ogawa, K.; Yamano, A.; Ohara, K.;
Yamaguchi, K.; Kawano S.; Tanaka, K.; Araki, Y.; Wada,
T.; *Otsuki, J. (2013). Double Helices of a
Pyridine-Appended Zinc Chlorophyll Derivative. *J. Am.
Chem. Soc.*, 135, 5262-5265.

2014

1. *Tominaga, M.; Yoneta, T.; Ohara, K.; Yamaguchi, K.; Itoh,
T.; Minamoto, C.; *Azumaya, I. (2014). Self-Assembly of a
Tetrapodal Adamantane with Carbazole Branches into
Hollow Spherical Aggregates in Organic Media.
Org. Lett., 16, 4622-4625.
2. *Tominaga, M.; Ukai, H.; Katagiri, K.; Ohara, K.;
Yamaguchi, K.; *Azumaya, I. (2014). Tubular Structures
Bearing Channels in Organic Crystals Composed of
Adamantane-Based Macrocycles. *Tetrahedron*, 70,
2576-2581.
3. *Tominaga, M.; Iekushi, A.; Katagiri, K.; Ohara, K.;
Yamaguchi, K.; *Azumaya, I. (2014). Channel-Dependent
Conformations of Single-Strand Polymers in Organic
Networks Composed of Tetrapodal Adamantanes with
N-heterocyclic Moieties. *Tetrahedron Lett.*, 55,
5789-5792.
4. *Tominaga, M.; Ohara, K.; Yamaguchi, K.; *Azumaya, I.
(2014). Hollow Sphere Formation from a
Three-Dimensional Structure Composed of an
Adamantane-Based Cage. *J. Org. Chem.*, 79, 6738-6742.
5. Shinozaki, Y.; Yoshikawa, I.; Araki, K.; Ohara, K.;
Yamaguchi, K.; Kawano, S.; Tanaka, K.; Araki, Y.; Wada,
T.; *Otsuki, J. (2014). Coordination Oligomers and
Polymers of an Oxazole-appended Zinc Chlorophyll
Derivative. *Chem. Lett.*, 43, 862-864.

2013

1. *Tominaga, M.; Iekushi, A.; Katagiri, K.; Ohara, K.;
*Yamaguchi, K. (2013). Organic Crystals Bearing Both
Channels and Cavities Formed from Tripodal Adamantane
Molecules. *Journal of Molecular Structure*, 1046, 52-56.
2. *Danjo, H.; Iwaso, K.; Kawahata, M.; Ohara, K.; Miyazawa, T.;
Yamaguchi, K. (2013). Preparation of
Tris(spiroorthocarbonate) Cyclophanes as Back to Back
Ditopic Hosts. *Org. Lett.*, 15(9), 2164-2167.
3. *Ohara, K.; Tominaga, M.; Azumaya, I.; *Yamaguchi, K.
(2013). Solvent-dependent Assembly of Discrete and
Continuous CoCl₂ Adamantane-based Ligand Complexes:
Observations by CSI-Mass Spectrometry. and X-ray
Crystallography. *Anal. Sci.*, 29(8), 773-776.



Laboratory for Neural Circuit Systems
Institute of Neuroscience

Staff

Takashi Tominaga, Ph.D.

Associate Professor since 2005

D.Sc. in University of Tsukuba, 1994

Yoko Tominaga

Research Assistant since 2006

Makiko Taketoshi

Research Assistant since 2016

Research

Since the expansion of the Institute of Neuroscience, Tokushima Research areas of the laboratory

I. Study of neural circuit mechanisms of learning and memory with optical recording methods

The primary interest of the laboratory is the neural circuit mechanisms of higher cognitive functions, such as learning and memory, in the brain. A measurement method that makes the laboratory unique in the field is an optical recording method that uses voltage-sensitive dye (VSD) with electrophysiology. As one of the leading laboratories in the use of this technique, we have been continuously developing the method since the 90s and have provided established tools to colleges throughout the world.

II. Analysis of the electrophysiological control of excitable membranes in connection with ciliary structures.

By focusing on the role of information integration in the membrane potentials of cells, we have used the model organism, paramecium, which is the simplest single-celled animal, to study the mechanisms of the membrane potential control of cilia.

Specific research aims

Area I

1. Neural circuitry mechanisms of the limbic system: Optical study. The limbic system is a brain structure that is critical for emotion and declarative memory. The limbic system consists of many areas, including the hippocampus, amygdala, and associated cortical systems, such as the entorhinal and piriform cortices. We are analyzing the function of these circuits by visualizing neural activity with the VSD optical recording methods.

We have revealed reverberation circuits and information integration mechanisms in the deep layers of the entorhinal and piriform cortices in association with the hippocampal neural circuit (Science, 1996; Neurosci. Res., 2008) with Professor Toshio Iijima's group at Tohoku University. In addition, we have found that neuronal signals from layer III of the medial entorhinal cortex

are critical for temporal association memory formation (Science, 2011) with Professor Susumu Tonegawa's laboratory at the Picower center for learning and memory at MIT. We have also revealed information integration processes in the entorhinal cortex (Eur. J. Neurosci., 2007) with Dr. Riichi Kajiwara and Dr. Ichiro Takashima's group at AIST Japan. We showed that the D-current plays an important role in the integration of neural activity in the entorhinal cortex in collaboration with Dr. Riichi Kajiwara (Japan Society for Neuroscience, 2012; Society for Neuroscience, 2012; supported by KAKENHI).

2. Development of an optical measurement microscope: stimulation pattern with a confocal microscope system and a new optical measurement.

The optical recording method with VSD requires high-speed and low-noise imaging. This requires new special optics. We have been developing special optics that meet these requirements (J. Neurosci. Methods, 2000; now commercially available as THT-microscope, BrainVision).

We have also developed special new ultra-high-speed and low-noise confocal optics (submitted; supported by JST tansaku, A-STEP). We have developed a microscope that allows us to conduct light stimulus patterns to the neural networks (SFN abstr., 2011). Recently, we have started a project to develop a special chamber that is suitable for these experiments (Supported by JST A-STEP, 2012-2013).

3. Mechanisms of late-onset brain dysfunctions caused by early transient exposure to chemicals and drugs.

There are several lines of evidence that indicate that the early transient exposure to certain chemicals during brain development results in the malfunctioning of cognitive function in adulthood. The neural mechanisms of these effects are largely unknown. We are evaluating these neural mechanisms with our optical recording methods as part of the research team that is supported by the Ministry of Health, Labour and Welfare (2008-, 2015-).

We have shown that the administration of valproic acid, which is the first-line drug used in the treatment of epilepsy, during pregnancy causes a collapse of the balance of excitation and inhibition in children born to these mothers (Japanese Society of Toxicology, 2012). This study is joint research that is being conducted with Kentaro Tanemura sensei of Tohoku University, Dr. Yoshikazu Nakajima of Nara Institute of Science and Technology, and the teacher Katsuhide Igarashi of the Japan Institute of Health Sciences. We organized a symposium at the Japan Neuroscience Society in Kyoto in June 2013.

4. Study of the Application of optical measurement methods to test

Laboratory for Neural Circuit Systems
Institute of Neuroscience

ES cell function.

This study was initiated in 2012 and is intended to use the optical recording method with VSD to characterize cells that are differentiated from human ES cells. This is joint research that is being conducted with Prof. Katsunori Sasaki, Shinshu University [supported by KAKENHI (A)].

5. Visualization of cell-specific membrane potential responses by the introduction of voltage sensitive fluorescent protein (VSFP), which is a new membrane potential-sensitive protein.

In collaboration with Dr. Thomas Knopfel at RIKEN BSI, we have succeeded in detecting optical signal-specific hippocampal pyramidal cells by introducing a new VSFP from 2012. The detection of cell-specific signals are made possible in specimens in vivo by the further development of this technique.

6. Detection and use of the optical signals from neural excitation with a polarized light microscope.

This is joint research that is being conducted with Dr. Tomomi Tani and Dr. Oldenburg of Woods Hole MBL. In this study, we aim to detect changes in nerve optical properties, such as polarization, that are caused by nerve excitation. In March of 2013, we will visit the MBL for this purpose.

7. Studies of the mechanisms of regulation by a variety of factors and the neural responses of hippocampal neural synapses.

We are collaborating with various laboratories to apply our method in order to examine the neural pathologies of diseases, such as Alzheimer's disease, and other factors (J. Neurosci., 1996; Neurosci. Letters, 1997; J. Neurosci., 2002; PNAS, 2004; Neuropharmacol., 2005).

8. Regulation mechanisms of neural activity by inhibitory synapses in the hippocampus.

The unique feature of VSD imaging compared to other biological imaging methods is that it can measure hyperpolarization and, thus, inhibitory neural responses. From this point of view, we found depolarizing GABA responses in area CA1 in response to tetanic stimulation (J. Neurophysiol., 2002; Pflüger's Arch., 2010). In addition, we found perisomatic inhibitory actions with feedforward inhibition (Neurosci. Res., 2009).

Area II

1. Physiological studies of osmoregulatory mechanisms and contractile vacuoles of Paramecium.

For the first time, we have applied electrophysiological methods to the study of the Paramecium organelles, the contractile vacuoles, and have revealed the membrane dynamics that are involved in this periodic activity (J. Exp. Biol., 1997a, b; 1998a, b; J. Cell Sci., 1999; J. Exp. Biol., 2005).

2. Physiological studies of membrane proteins and cilia of paramecium response mechanisms.

The use of recent techniques of RNA interference knockdown in

combination with the whole genome project of the Paramecium has enabled us to knock down particular proteins that are associated with cilia disease (so-called ciliopathy). We have found that the absence of some molecules that have been thought to be structural proteins induces behavioral changes. By applying electrophysiological methods to this mutant, we have examined the relationship of that behavior and the membrane responses and found that some of these "structural proteins" are actually involved in membrane potential-mediated signal transduction (e.g., Eukary. Cell, 2012). This is joint research that is being conducted with Prof. Hori of Yamaguchi University.

Publications (2012~2017)

[Original papers]

*Corresponding author

2016

1. Yoshimura, H., Sugai, T., Kato, N., Tominaga, T., Tominaga, Y., Hasegawa, T., Yao, C., and Akamatsu, T. Interplay between non-NMDA and NMDA receptor activation during oscillatory wave propagation: Analyses of caffeine-induced oscillations in the visual cortex of rats. *Neural Networks*. 79:141-149 (2016) DOI: 10.1016/j.neunet.2016.03.012
2. *Tominaga T and Tominaga Y (2016). Paired burst stimulation causes GABAA receptor-dependent spike firing facilitation in CA1 of rat hippocampal slices. *Front. Cell. Neurosci.* **10**:9. doi: 10.3389/fncel.2016.00009

2015

3. Juliandi B, Tanemura K, Igarashi K, Tominaga T, Furukawa Y, Otsuka M, Moriyama N, Ikegami D, Abematsu M, Sanosaka T, Tsujimura K, Narita M, Kanno J, Nakashima K (2015) Reduced Adult Hippocampal Neurogenesis and Cognitive Impairments following Prenatal Treatment of the Antiepileptic Drug Valproic Acid. *Stem Cell Rep.*:1-14.. doi:10.1016/j.stemcr.2015.10.012

2013

4. Takashi Tominaga^{CA}, Riichi Kajiwara, and Yoko Tominaga (2013) VSD imaging method of ex vivo brain preparation *Journal of Neuroscience and Neuroengineering* 2, 211-219 (2013) [Featured Article に採用]
5. Tominaga T^{CA} and Tominaga Y. (2013) A new non-scanning confocal microscopy module for functional voltage-sensitive dye and Ca²⁺ imaging of neuronal circuit activity *Journal of Neurophysiology* J Neurophysiol 110, 553-561; published ahead of print April 24, [Also featured as Key Scientific Articles on Global Medical Discovery]